

Older Adults and Falls

Joe Rosenthal, MD, MPH
Assistant Professor, Clinical
Department of Physical Medicine & Rehabilitation
The Ohio State University Wexner Medical Center


Objectives

- Introduce the epidemic of Older Adult Falls
- Discuss risk factors for falling
- Discuss interventions and resources available

Falls are not a normal part of aging, they can be prevented

How bad is the Problem

- One out of three older adults falls each year
- Each year, 2.5 million older people are treated in emergency departments for fall injuries.
- 700,000+ patients a year are hospitalized because of a fall injury, most often because of a head injury or hip fracture (250,000 people).



<http://www.cdc.gov/injury/wisqars/>
3/2016

Falls Are Serious and Cause Injuries

- 20% of falls cause a serious injury such as broken bones or a head injury.
- Almost all hip fractures are caused by falling.
- Falls are the most common cause of traumatic brain injuries (TBI).

CDC website
3/2016

Falls are costly

- Direct medical costs for fall injuries are \$34 billion annually.
- Hospital costs account for two-thirds of the total.
 - The average hospital cost for a fall injury is \$35,000.
- Fall injuries are among the 20 most expensive medical conditions.
- The costs of treating fall injuries goes up with age.
- Medicare pays for about 78% of the costs of falls

CDC website
3/2016

Major Problem (Epidemic) in Ohio

Falls are the #1 cause of injuries leading to ER visits, hospitalizations and deaths for Ohioans age 65+:

- An injury every 2.5 minutes
- An emergency room visit every 8 minutes
- Two hospitalizations each hour
- Three deaths each day

Direct care costs alone = \$646 million

Ohio Department of Health

Major Problem (Epidemic) in Ohio

- Ohioans age 65+ make up approximately 14% of the population, but account for more than 83% of fatal falls.
- Fatal falls among older Ohioans increased 167% from since 2000.
- 1 in 3 Ohioans age 65+ living in the community fall each year. 1 in 2 after age 79.
- More than half of older adults who live in a nursing home will fall this year.

Ohio Department of Health

Why isn't more being done?

- Clinical assessment and intervention by healthcare providers can significantly reduce falls.
 - The literature also supports screening and management of falls in primary care settings.
 - Guidelines have been developed by the American Geriatrics Society that suggest at least yearly screening for falls in older adults.
- Despite these recommendations, patients often will not speak with their healthcare providers about falling.
 - Less than half of older adults who fall discuss it with their healthcare provider.

Why aren't falls discussed?

- Reasons providers are not discussing with patients are many:
 - not recognizing the problem
 - not being aware of resources available to help
 - not having time to implement into practice
- Older adults may be hesitant to discuss falls with their providers:
 - fear of losing their independence
 - not knowing that steps can be done to reduce the risk of falling

Risk Factors for Falling

Risk Factors for Falling

- Prior Falls
- Gait disturbance and impaired balance
- Inactivity and/or generalized weakness
- Polypharmacy and certain medication classes
- Chronic pain
- Impaired vision and hearing
- Impaired cognition
- Impaired sensation/proprioception
- Dizziness and vertigo
- Fear of falling
- Depression and other psychiatric disorders
- Environment

Fear of Falling

- Fear of falling is highly associated with future falls
- Just asking a person if they are afraid of falling goes a long way to predicting future falls
 - Is it situational?
 - Has it caused lifestyle changes or decreased quality of life?

Gait Disturbance

- Short shuffling steps
- Wide based support
- Improper use of cane or walker
 - Including not using

Balance Disturbance

- Easily perturbed
- Balance worsens if looking in a different direction or in the dark
- Feeling lightheaded or dizzy

Polypharmacy and Falls

- Being on multiple medications is associated with falls
 - Indicator of multiple disease processes that increase fall risk
 - Drug interactions, side effects of the medications
- No specific cutoff, but studies often use 5 or 6 and greater

Medication Drug Classes that Increase Fall Risk

- Benzodiazepine
- Narcotics
- Anti-depressants
- Diabetic medications
- Blood pressure medications
- Sleep aide medications

Dizziness/vertigo/postural hypotension

- Multiple etiologies, but not uncommon in elderly
- Inner ear problem
 - Menier's Disease
 - Vestibular neuritis
 - BPPV
- Vision changes
- Hypotension
- Medication side effect
- Near-syncope/syncope

Sensory Changes

- Impaired vision
- Impaired hearing
- Decreased sensation/proprioception

Cognition and Falls

20

How prevalent is cognitive deficits in older adult fallers?

- Fallers with femoral neck fracture:
 - 49% had difficulties with orientation on MMSE
 - 70% had difficulties with at least some short term memory deficits as seen on MMSE

Tsur et al 2014

Cognitive Deficits, Aging, & Falls

- Falls are associated with cognitive deficits, even if criteria for Mild Cognitive Impairment or dementia are not met
 - Neuropsychological assessments have shown relationship between falls and cognitive deficits associated with normal aging.
 - Lower scores of processing speed and executive function associated with single falls and recurrent falls
- Studies have shown that all levels of cognitive impairment (aging to dementia) are associated with higher fall rates with dementia having highest fall risk.

Cognition Components That Pertain to Fall Risk

- Executive Function: term used to describe brain's management of multiple cognitive processes
 - Planning
 - Judgment
 - Impulsivity
- Attention
 - Divided Attention and Dual-Task
- Processing Speed
- Memory
 - Working Memory: ability to take information and process it instantaneously (different than short term memory)

What can be done?

24

Resources/Guidelines

- Ohio Department of Health: Steady U
 - <http://aging.ohio.gov/steadyu/>
- CDC: STEADI
 - <http://www.cdc.gov/homeandrecrereationalsafety/falls/>
- American Geriatrics Society
 - <http://geriatricscareonline.org/toc>

Non-Pharmacologic Interventions

- Ask about and screen for risk factors, including cognition
- Educate
- DME
 - Be mindful of dangerous situations with cog deficits using DME (i.e. scooters/power wheelchairs)
- Assistive technology
 - Bed alarms, door alarms, lights with motion sensors
- Therapy (PT, OT, ST)
- Daily exercise programs
- Refer to specialists and/or further testing if warranted (tilt table, EMG, etc)

Pharmacologic Management

- Reduce cognitive slowing and sedating meds
 - benzodiazepines
 - pain medications
 - sleep medications
 - psychotropic
- Limited research on specific cognitive medications that reduce fall risk
 - Stimulants
 - Memory/Executive Function Enhancers

Vitamin D Supplementation

- Studies have shown that supplementation with Vitamin D (even if not deficient) can reduce fall rates
 - Typically 800 IU/day are recommended

American Geriatrics Society Workshop 2014

Environmental Hazards

Potential Hazards in home environment that can cause falls

- Throw rugs
- No grab bars
- Showers without anti-slip surfaces
- Lack of handrails
- Small animals
- Clutter
- Unlit areas
- Overhead lights that are not easily reached
- Damaged floors

Other Interventions

- Proper footwear
- Yearly eye exams
- Screen for and treat hearing loss

What Can be Done to Improve Safety?

- Discussion between healthcare provider and patient
- Look for community resources
- Assess home environment for hazards and address
 - Consider Home Assessment by OT or other healthcare provider
- Encourage use cane or walker if they have been recommended
- Ensure medications are being taken appropriately, monitor for side effects, and review medication lists so that those no longer needed can be discontinued
- Encourage activity and exercise

Refer to Therapy

- Therapies (PT/OT) can help reduce risk of falls for those at risk
 - Improved strength and balance
 - Proper use of DME
 - Improved independence with ADLs
 - Home assessments

Refer to Community Based Exercise Programs

- Literature supports exercise programs that reduce falls
 - Tai Chi
 - Matter of Balance
 - Otago exercise program

References

- Auriel E, Hausdorff JM, Herman T, et al. Effects of methylphenidate on cognitive function and gait in patients with Parkinson's disease: a pilot study. *Clin Neuropharmacol.* 2006;29:15-17.
- American Geriatrics Society Workgroup on Vitamin D Supplementation for Older Adults. Recommendations abstracted from the American Geriatrics Society consensus statement on Vitamin D for prevention of falls and their consequences. *J Am Geriatr Soc.* 2014; 62(1): 147-152.
- Ben Itzhak R, Giladi N, Gruendlinger L, et al. Can methylphenidate reduce fall risk in community-living older adults? A double-blind, single-dose cross-over study. *J Am Geriatr Soc.* 2008;56:695-700.
- Chung KA, Lobb BM, Nutt JG, et al. Effects of a central cholinesterase inhibitor on reducing falls in Parkinson disease. *Neurology.* 2010;75:1263-1269.
- Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, Lamb SE. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev.* 2012 Sep 12;9:CD007146. doi: 10.1002/14651858.CD007146.pub3. Review.
- Gleason CE, Gangnon RE, Fischer BL, Mahoney JE. Increased Risk for Falling Associated with Subtle Cognitive Impairment: Secondary Analysis of a Randomized Clinical Trial. *Dement Geriatr Cogn Disord.* 2009 Jul; 27(6): 557-563. doi: 10.1159/000228257
- Hebert LE, Weuve J, Scherr PA, Evans DL. Alzheimer disease in the United States (2010-2050) estimated using the 2010 census. *Neurology.* 2013;80:1778-83.

References

- Herman T, Mirelman A, Giladi N, Schweiger A, Hausdorff JM. Executive control deficits as a prodrome to falls in healthy older adults: a prospective study linking thinking, walking, and falling. *J Gerontol A Biol Sci Med Sci.* 2010 Oct;65(10):1086-92. doi: 10.1093/gerona/gdq077.
- Mirelman A, Herman T, Brozgol M, Dorfman M, Sprecher E, Schweiger A, Giladi N, Hausdorff JM. Executive function and falls in older adults: new findings from a five-year prospective study link fall risk to cognition. *PLoS One.* 2012;7(6):e40297. doi: 10.1371/journal.pone.0040297.
- Montero-Odasso M, Muir-Hunter SW, Oteng-Amoako A, et al. Donepezil improves gait performance in older adults with mild Alzheimer's Disease: a phase II clinical trial. *Journal of Alzheimer's Disease* 2015; 43(1):193-9. doi: 10.3233/JAD-140759
- Moyer VA; U.S. Preventive Services Task Force. Prevention of falls in community-dwelling older adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2012 Aug 7;157(3):197-204.
- O'Halloran AM, Penard N, Galli A, Wei Fan, C, Robertson IH, Kenny RA. Falls and falls efficacy: the role of sustained attention in older adults. *BMC Geriatrics* 2011; 11(85). doi:10.1186/1471-2318-11-85.
- Holtzer R, Friedman R, Lipton RB, Katz M, Xue X, Verghese J. The relationship between specific cognitive functions and falls in aging. *Neuropsychology.* 2007 September ; 21(5): 540-548. doi:10.1037/0894-4105.21.5.540
- Phelan EA, Mahoney JE, Voit JC, Stevens JA. Assessment and Management of Fall Risk in Primary Care Settings. *Med Clin North Am.* 2015 Mar;99(2):281-293. doi: 10.1016/j.mcna.2014.11.004. Review.

References

- Smith-Ray, R.L., Hughes, S.L., Prohaska, T.R., Little, D.M., Jurivich, D.A. & Hedeker, D. (2013) Impact of cognitive training on balance and gait in older adults. *Journals of Gerontology, Series B: Psychological Sciences and Social Sciences*, 70(3), 357–366, doi:10.1093/geronb/gbt097.
- Stevens JA, Ballesteros MF, Mack KA, Rudd RA, DeCaro E, Adler G. Gender differences in seeking care for falls in the aged Medicare population. *Am J Prev Med*. 2012 Jul;43(1):59-62. doi: 10.1016/j.amepre.2012.03.008.
- Tsur A, Eluz D, Itah D, Segal Z, Shakeer N, Galin A. Clinical profile of fallers with femoral neck fractures. *PM&R* 2014; 6:390-394. doi: 10.1016/j.pmrj.2013.10.013.
- Ward A, Arrighi HM, Michels S, Cedarbaum JM. Mild cognitive impairment: disparity of incidence and prevalence of estimates. *Alzheimer's & Dementia* 2012; 8(1): 14-21. doi:10.1016/j.jalz.2011.01.002

Prescribing walkers and canes

When to consult the physical medicine physician

Fall Prevention in Order Adults: Therapy Perspective

ReNea L. Owens, PT
Clinical Manager
Outpatient Rehabilitation
The Ohio State University Wexner Medical Center

Objectives

- Introduce evidenced based tests and screening measures to identify fall risk
- Identify treatment interventions to decrease fall risk
- Identify community resources to elderly fall prevention programs



Evidenced Based Tests and Screening Methods to Assess Fall Risk

- 30 second sit to stand
- 4 Stage Balance Test
- Timed up and Go (TUG)
- Functional Reach Test
- 5 times sit to stand
- Functional Gait Assessment
- 4 Square Step Test

STEADI

STEADI – Stopping Elderly Accidents, Deaths & Injuries

- Center for Disease Control and Prevention (CDC) - a tool kit for physician offices to assess risk
- 30 Second Chair Stand Test
- 4 Stage Balance Test
- Timed Up and Go (TUG)



Don't Fall Behind on Fall Prevention.

www.cdc.gov

STEADI manages risks and saves lives.



1 in 4 older adults will fall this year.

www.cdc.gov

Keep seniors STEADI by screening for fall risk.

<http://www.cdc.gov/steady>

Fall Risk Assessment

30-Second Chair Stand Test
Assesses leg strength and endurance



Fall Risk Assessment

30 Second Chair Stand Test

Results

30 Second Chair Stand (# Stands)

Age	Men			Women			
	• Below Average	Average	Above Average	Age	Below Average	Average	Above Average
60-64	<14	14-19	>19	60-64	<12	12-17	>17
65-69	<12	12-18	>18	65-69	<11	11-16	>16
70-74	<12	12-17	>17	70-74	<10	10-15	>15
75-79	<11	11-17	>17	75-79	<10	10-15	>15
80-84	<10	10-15	>15	80-84	<9	9-14	>14
85-89	<8	8-14	>14	85-89	<8	8-13	>13
90-94	<7	7-12	>12	90-94	<4	4-11	>11

4-Stage Balance Test

feet together semi-tandem full tandem



Timed Up and Go (TUG)

- Assess Mobility
- Assess Postural Stability, Gait, Stride length, Sway
- Without a device greater than 12.0 sec fall risk
- With a device greater than 20 seconds fall risk




A ————— B



Functional Gait Assessment

- Assesses postural stability during various walking tasks for 20 feet
- 10 item task
- Score 0-3 with a total of 30 points
- < 22/30 Predicts Falls
- < 21/30 Falls in next 6 months



5 Times Sit to Stand

- Assesses lower limb functional strength
- 15 second cut off for fallers

- 11.4 Norms for 60s
- 12.6 Norms for 70s
- 14.8 Norms for 80s



Functional Reach Test

- Assesses stability with maximum reach from a standing fixed position


- Less than 6 inches Hi Risk of Falls
- 6-10 inches Mod Risk of Falls
- Greater than 10 inches Low risk of Falls



4 Square Step Test

- Assesses dynamic balance forward, backwards, and lateral movements stepping over crossed canes

- Greater than 15 seconds at risk for Multiple falls
- 10-14 seconds moderate Fall Risk
- Less than 10 seconds no fall risk



Interventions

Otago Exercise Program

Flexibility



Balance Exercises



Strengthening



Walking



Source: Some images courtesy of NIA

Community Resources

www.cdc.gov/steady

Stopping Elderly Accidents, Deaths & Injuries

- Information for health care providers for fall risk assessments
- Education materials to assist patients

Community Resources

Matter of Balance

8-week structured group intervention

Strategies to reduce fear of falling

Increase activity levels

Change their environment to reduce fall risk

Exercise to increase strength and balance



Community Resources

Tai Chi for Arthritis

Tai Chi one of the most effective exercises for preventing falls.

Improve all muscular strength, flexibility, balance, stamina, and more.



Community Resources

Stepping On

- strategies, and exercises to reduce falls
- increase self-confidence in situations that they are at risk of falling



Community Resources

Stay Active and Independent for Life (SAIL)

A strength, balance and fitness program 3 times a week in a one hour class, in sitting or standing. Community-dwelling older adults (65+) and people with a history of falls.



References

1. Centers for Disease Control and Prevention, National Center for Injury Prevention and Control, Division of Unintentional Injury Prevention
2. Wrisley DM & Kumar NA, Functional Gait Assessment: Concurrent, Discriminative, and Predictive Validity in community-Dwelling Older Adults, Physical Therapy, May 2010, vol 90 (5): 761-773
3. Criter, RE & Honaker JA, Identifying Balance Measures Most Likely to Identify Recent Falls, The Journal of Geriatric Physical Therapy, Jan-Mar 2016, 39 (1):30-37

References

4. Muir SW et al, The Role of Cognitive Impairment in Fall Risk among Older Adults: A systematic Review and Meta-Analysis, Age-Aging, 2012 41 (3): 299-308
5. Dorfman M et al, Dual-Task Training on a Treadmill to Improve Gait and Cognitive Function in Elderly Idiopathic Fallers, Journal of Neurology Physical Therapy, Oct 2014, 38:246-253
6. National Council on Aging (NCOA)
<https://www.ncoa.org/healthy-aging/falls-prevention/falls-prevention-programs-for-older-adults/>

References

7. Lee J et al, Analytical Review: Focus on Fall Screening Assessments, The American Academy of Physical Medicine and Rehabilitation, July 2013, 5:609-621
8. Rehabilitation Measures Database
<http://www.rehabmeasures.org/default.aspx>