

Common Fractures

Ryan K. Harrison, MD
Orthopaedic Trauma
Assistant Professor
Orthopaedic Surgery
The Ohio State University Wexner Medical Center

Objectives

- **Identify Common Fractures**
- **Discuss initial treatment of fractures**
- **Discuss definitive treatment of common fractures and expected outcomes**

Statistics

- **About 18.5 million visits to physicians per year are for fractures**
- **The majority of these occur in the outpatient setting**
- **MSK injuries result in loss of function and negatively impact our economy**
- **Our aging population will result in increased fracture – especially fragility fractures – with even more loss of function**

Most Common Fractures

- **Clavicle**
- **Ankle**
- **Pediatric Forearm Fractures**
- **Wrist**
- **Hip**

Initial Evaluation

- **Identify injured extremity**
- **Is there deformity?**
- **Neurovascular Evaluation**
- **Skin integrity**
- **Radiographs**

Radiographic Principles

- **Image the injured bone**
- **Obtain radiographs of the joint ABOVE and BELOW the injured bone**

Initial Treatment

- Immobilize with splint
- Elevate injured extremity to lessen swelling
- Ice
- Anti-inflammatory medication for pain/swelling
- Opioids for severe pain
- Refer for definitive management

General Treatment Principles

- Non-displaced and stable fractures can be treated with non-operative management
- Displaced, open and unstable fractures should be treated surgically
- Encourage early range of motion when able
- Weight bearing is limited for minimal necessary time
- Immobilization may be necessary for appropriate healing

Clavicle Fractures

- **Common in falls onto an extremity**
- **Non-operative management with sling has been the traditional approach**
- **A large number can still be treated non-operatively**



- Recent research points more towards operative management for severely displaced fractures



- **Regardless of treatments – patients are treated in a sling for 6-8 weeks**
- **Weight bearing as tolerated**
- **Range of motion above shoulder level is restricted for 4-6 weeks**
- **Full weight bearing and strengthening is permitted after about 3 months**

Ankle Fracture

- **Twisting injury leads to predictable injury pattern**
- **Fibula fractures alone can often be treated non-operatively – decision depends on stability of the ankle joint**
- **Bimalleolar and Trimalleolar fractures often require operative fixation**



Stress View Radiograph





- **Stable fractures can be treated without surgery**
 - **Early weight bearing**
 - **Edema Control**
 - **RICE**
 - **RANGE OF MOTION**
- **Bone healing takes 6-8 weeks, but full recovery can be 3-6 months**

- **Unstable injuries require operative fixation**
- **Non-weight bearing after operative fixation for 6-12 weeks depending on the severity of the injury**
- **Encourage ROM after early immobilization period**
- **Full recovery can be 6-12 months**

Special Populations

- **Diabetic patients require more aggressive treatment and longer non-weight bearing**
- **Obese patients should be considered for longer non-weight bearing**
- **Skin condition in geriatric patients should be carefully monitored**

Pediatric Forearm/Wrist Fractures

- **Pediatric forearm and wrist fractures are among the most common fractures in children under 14**
- **These fractures are often caused by a fall from a height – monkeybars and trampolines**
- **Open physes allow for non-operative management of these fractures in most circumstances**
- **Closed reduction and casting is the most common method of treatment**



Source: eORIF.com

- **6-8 weeks of cast treatment followed by removable splint for an additional 4-6 weeks**
- **After cast removal – encourage range of motion and slow return to activity**
- **Open physes allow for remodeling of fractures and differing acceptable angles of reduction depending on patient age**
- **Older patients require more perfect reductions and may require surgical fixation**

Fragility Fractures

- **8.9 million fractures worldwide related to osteoporosis**
- **Hip, wrist and vertebral fractures occur in nearly equal numbers**
- **1:3 women over 50 incur fractures**
- **1:5 men over 50 incur fractures**
- **These numbers are expected to increase dramatically as our population ages**

Distal Radius Fracture

- **Often caused by fall on outstretched hand**
- **Early treatment includes reduction and splinting**
- **Non-surgical treatment can be acceptable in cases of appropriate alignment after reduction and splinting with transition to casting**
- **Surgical treatment is indicated for younger, active patients and in those with unacceptable reduction**



- **In non-operative management, cast for 6-8 weeks, followed by brace wear and focus on wrist range of motion**
- **Even with significant deformity – function can be appropriately regained in low demand individuals**

- **Operative fixation allows for early range of motion**
- **Bone healing can take 6-8 weeks**
- **Return to function is expected with appropriate treatment**

Hip Fracture

- **75% hip fractures occur in women**
- **10-20% of patients who were community ambulators prior to their injury lose their ability to function independently**
- **Current incidence around 1.6million/year – expected to increase to as much as 6 million/ year by 2050**
- **20-25% of patient die within one year after sustaining a hip fracture**

- **Much research is focused on improving outcomes after hip fracture**
- **Identification of at-risk patients and measures to protect bone health**
- **Earliest possible treatment of appropriately risk stratified patients may lead to improved mortality and preservation of pre-injury functional status**
 - **This is a subject of ongoing research at OSU participating in an international study**

- **Patients are treated in a fashion similar to acute coronary patients with dedicated teams and surgical fixation of their fractures in an expedited fashion**

- **Risk/benefit analyses are always necessary when considering surgical procedures and not every patient should undergo emergent procedures.**
- **Some patients with acute medical conditions benefit from medical optimization prior to surgery.**
- **A minority of patients are not surgical candidates and palliative measures should be discussed with the patient and family.**

- **Patients who have never undergone osteoporosis workup should have workup initiated by hospital medical/orthopaedic personnel**
 - **Laboratory workup for secondary osteoporosis**
 - **DEXA**
 - **Geriatric Consultation**
 - **Endocrine evaluation**
- **Follow-up after hospitalization should be arranged with appropriate personnel to initiate appropriate treatment**

- **Goals of surgical treatment are to mobilize patients quickly with full weight bearing**
 - **Options are fixation of the fracture or arthroplasty procedure**
- **Fixation is reserved for non-displaced or younger patients with displaced fractures**
- **Partial arthroplasty is indicated in low demand individuals**
- **Total hip arthroplasty is indicated in younger, more active individuals**





