

Hypertension Update

Aaron J. Friedberg, MD
Assistant Professor, Clinical
Division of General Internal Medicine
The Ohio State University Wexner Medical Center

Overview

- Background
- Diagnosis
- Management
- Guideline Comparison

Background

- Importance
 - ✓ In 2010, hypertension was the leading cause of death and disability-adjusted life years worldwide.
 - ✓ In the US, second only to cigarettes as preventable cause of death
 - ✓ 25% of cardiovascular events attributable to hypertension in ARIC

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Background

- Importance
 - ✓ Half with hypertension do not have adequate control

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Background

- **Prevalence**
 - ✓ 32% under JNC-8/ current ACP/AAFP
 - ✓ 46% under recent AHA/ACC

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Kovell LC et al. US Hypertension Management Guidelines: A Review of the recent Past and Recommendations for the Future. Journal of the American Heart Association. 2015;4:e002315, originally published December 7, 2015

Background

- **Complex & Interdependent relationship between modifiable & fixed risk factors**
- **Modifiable**
 - ✓ Cigarette smoke exposure
 - ✓ Diabetes
 - ✓ Dyslipidemia
 - ✓ Obesity
 - ✓ Lack of physical activity
 - ✓ Poor diet
- **Fixed**
 - ✓ Male
 - ✓ Family History
 - ✓ Low SES
 - ✓ OSA
 - ✓ Stress
 - ✓ Chronic Kidney disease
 - ✓ Age

Berry JD, Dyer A, Cai X, et al. Lifetime risks of cardiovascular disease. N Engl J Med. 2012;366:321-9

Diagnosis

- **Proper Blood Pressure Assessment**
 - ✓ 1: Patient preparation

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Proper Blood Pressure Assessment**
 - ✓ 1: Patient preparation
 - ✓ 2: Proper technique

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Proper Blood Pressure Assessment**
 - ✓ 1: Patient preparation
 - ✓ 2: Proper technique
 - ✓ 3: Proper measurements

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Proper Blood Pressure Assessment**
 - ✓ 1: Patient preparation
 - ✓ 2: Proper technique
 - ✓ 3: Proper measurements
 - ✓ 4: Good documentation

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Proper Blood Pressure Assessment**
 - ✓ 1: Patient preparation
 - ✓ 2: Proper technique
 - ✓ 3: Proper measurements
 - ✓ 4: Good documentation
 - ✓ 5: Average readings

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Proper Blood Pressure Assessment**
 - ✓ 1: Patient preparation
 - ✓ 2: Proper technique
 - ✓ 3: Proper measurements
 - ✓ 4: Good documentation
 - ✓ 5: Average readings
 - ✓ 6: Give patient the numbers

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Definitions (AHA/ACC):**
 - ✓ ->Normal <120/80mmHg
 - ✓ ->Elevated blood pressure: 120-129mmHg/<80mmHg
 - ✓ ->Stage 1 HTN: 130-139mmHg or diastolic 80-89mmHg
 - ✓ ->Stage 2 HTN: 140mmHg or greater systolic, 90mmHg or greater diastolic

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Other methods:**
- **Ambulatory Blood Pressure Monitoring**
 - ✓ ->24-hour mean of 125/75mmHg or more
 - ✓ ->Awake mean of 130/80mmHg or more
 - ✓ ->Asleep mean of 110/65mmHg or more

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Siu AL. Screening for high blood pressure in adults: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med. 2015;163:778-86.

Agarwal R, Bills JE, Hecht TJW, et al. Role of home blood pressure monitoring in overcoming therapeutic inertia and improving hypertension control: a systematic review and meta-analysis. Hypertension. 2011;57:29-38.

Diagnosis

- **Other methods:**
- **Ambulatory Blood Pressure Monitoring**
 - ✓ ->24-hour mean of 125/75mmHg or more
 - ✓ ->Awake mean of 130/80mmHg or more
 - ✓ ->Asleep mean of 110/65mmHg or more
- **Home Blood Pressure Monitoring**

Pickering TG, Miller NH, Ogedegbe G, et al. Call to action on use and reimbursement for home blood pressure monitoring: a joint scientific statement from the American Heart Association, American Society of Hypertension, and Preventive Cardiovascular Nurses Association. Hypertension. 2008;52:10-29.

Diagnosis

- **“What’s more important, the top number or the bottom number?”**
 - ✓ Under age 50: diastolic matters more

Taylor BC et al. Impact of diastolic and systolic blood pressure on mortality: implications for the definition of “normal.” J Gen Intern Med. 2011 Jul;26(7):685-90.

Diagnosis

- “What’s more important, the top number or the bottom number?”
 - ✓ Under age 50: diastolic matters more
 - ✓ Over age 50: systolic/pulse pressure is greater predictor of events

Taylor BC et al. Impact of diastolic and systolic blood pressure on mortality: implications for the definition of “normal.” J Gen Intern Med. 2011 Jul;26(7):685-90.

Franklin SS et al. Does the relation of blood pressure to coronary heart diseases risk change with aging? The Framingham Heart Study. Circulation. 2001;103(9):1245.

Diagnosis

- Clinical Evaluation-History
 - ✓ Organ damage
 - ✓ Risk factors
 - ✓ Lifestyle
 - ✓ Medications
 - ✓ Substance abuse

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- Clinical Evaluation-Physical Exam
 - ✓ End-organ damage
 - ✓ Fundoscopic exam

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- Clinical Evaluation-Lab Testing
 - ✓ Electrolytes with calcium and creatinine
 - ✓ Fasting glucose
 - ✓ Urinalysis
 - ✓ Complete Blood Count
 - ✓ Thyroid testing-TSH
 - ✓ Lipids
 - ✓ Electrocardiogram
 - ✓ Use labs & clinic data to calculate 10-year ASCVD risk (ACC/AHA)

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Chobanian AV et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. JAMA. 2003;289(19):2560.

Diagnosis

- **Clinical Evaluation-Lab Testing**
 - ✓ Additional depending on circumstance:
 - Urine albumin to creatinine ratio
 - Echocardiogram
 - Secondary Hypertension

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Chobanian AV et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. JAMA. 2003;289(19):2560.

Diagnosis

- **Risk Factors**
 - ✓ Primary
 - ✓ Secondary

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Diagnosis

- **Primary Risk Factors**
 - ✓ Family history-about 30% of population variation is due to genetic factors, and twice as common if at least 1 parent has it
 - ✓ Age
 - ✓ Sodium intake
 - ✓ Obesity
 - ✓ Physical activity level
 - ✓ Race
 - ✓ Alcohol
 - ✓ Kidney mass

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
Staessen JA et al. Essential Hypertension. Lancet. 2003;361(9369):1629.

Diagnosis

- **Secondary Risk Factors**
 - Oral contraceptives
 - NSAIDs
 - Antidepressants
 - Corticosteroids
 - Decongestants
 - Weight-loss medications
 - Antacids with sodium
 - Erythropoietin
 - Cyclosporine/tacrolimus
 - Stimulants
 - Atypical antipsychotics
 - Anti-angiogenesis
 - Tyrosine Kinase inhibitors
 - Illegal drugs
 - Kidney disease
 - Hyperaldosteronism
 - Renovascular
 - Obstructive Sleep Apnea
 - Pheochromocytoma
 - Cushing's syndrome
 - Endocrine disorders
 - Coarctation of the aorta

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
UpToDate.com. Overview of Hypertension in adults. Retrieved April 4, 2018.

Diagnosis

- **Complications**
 - ✓ Left ventricular hypertrophy
 - ✓ Heart Failure
 - ✓ Stroke-ischemic & hemorrhagic
 - ✓ Coronary Artery Disease
 - ✓ Chronic Kidney Disease/End-Stage Renal Disease

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017. UpToDate.com. Overview of Hypertension in adults. Retrieved April 4, 2018.

Management

- **Lifestyle**
- **Medications**

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Management

- **Lifestyle**
 - ✓ Salt restriction--5mmHg systolic/2.5mmHg diastolic
 - ✓ Increased potassium intake via diet
 - ✓ Weight loss--1mmHg/lb lost
 - ✓ DASH diet: combination of low-salt, high potassium, magnesium, calcium, protein, fiber, low fat/cholesterol, reduced by 6mmHg systolic/4mmHg diastolic
 - ✓ Exercise: 3-4x40minutes for 12 weeks can get 4-6/3mmHg
 - ✓ Limit daily alcohol intake-2 or fewer for men, 1 or fewer for women

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017. Eckel RH et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2014;63(25 Pt B):2960.

Management

- **Medication**
 - ✓ Effective on outcomes

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Management

- Medication-where to start?

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Management

- Medication-where to start?
 - ✓ Thiazide diuretics
 - ✓ Long-acting calcium channel blockers eg amlodipine (CCB)
 - ✓ Angiotensin converting enzyme inhibitors (ACE-I)
 - ✓ Angiotensin receptor blockers (ARB)

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Management

- Special cases:
 - ✓ Black patients-best evidence for starting with thiazide or CCB
 - ✓ Diabetic nephropathy or any chronic kidney disease with proteinuria-ACE-I/ARB
 - ✓ Don't start with a beta blocker anymore, unless they have coronary artery disease or systolic heart failure

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Management

- Special cases:
 - ✓ Ischemic Heart Disease
 - ✓ Heart Failure
 - ✓ Chronic Kidney Disease
 - ✓ Renal Transplant
 - ✓ Secondary Stroke Prevention
 - ✓ Pregnancy
 - ✓ Hypertensive crisis
 - ✓ Cognitive Decline
 - ✓ Sexual Side Effects
 - ✓ Major surgery

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Management

- Sequential Management
 - ✓ 15mmHg+ above goal: start with 2 agents
 - ✓ 2 is not enough: go to 3 (ACE-I/ARB, thiazide, CCB)
 - ✓ 3 is not enough: “drug-resistant hypertension)
 - ✓ 4 drugs+: resistant hypertension

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Guideline Comparison: Definition

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>	<u>Stage II</u>	
<u>>140/90mmHg</u>	<u>Stage II</u>	
<u>130-139/80-89mmHg</u>	<u>Stage I</u>	
<u>120-129/<80mmHg</u>	<u>Elevated</u>	
<u><120/80mmHg</u>	<u>Normal</u>	

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Guideline Comparison: Definition

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>	<u>Stage II</u>	
<u>>140/90mmHg</u>	<u>Stage II</u>	
<u>130-139/80-89mmHg</u>	<u>Stage I</u>	<u>Does not specifically define hypertension/prehypertension</u>
<u>120-129/<80mmHg</u>	<u>Elevated</u>	
<u><120/80mmHg</u>	<u>Normal</u>	

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.

Guideline Comparison: Treatment

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>		
<u>>140/90mmHg</u>		
<u>130-139/80-89mmHg</u>		
<u>120-129/<80mmHg</u>		
<u><120/80mmHg</u>		

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APha/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
Gaseem et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinical Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. Ann Intern Med. 2017;166(6):430-437.

Guideline Comparison: Treatment

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>		
<u>>140/90mmHg</u>		
<u>130-139/80-89mmHg</u>		
<u>120-129/<80mmHg</u>		
<u><120/80mmHg</u>	<u>Screening</u>	<u>Screening</u>

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
Gaseem et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinician Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. Ann Intern Med. 2017;166(6):430-437.

Guideline Comparison: Treatment

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>		
<u>>140/90mmHg</u>		
<u>130-139/80-89mmHg</u>		
<u>120-129/<80mmHg</u>	<u>Close monitoring</u>	<u>Close monitoring</u>
<u><120/80mmHg</u>	<u>Screening</u>	<u>Screening</u>

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
Gaseem et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinician Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. Ann Intern Med. 2017;166(6):430-437.

Guideline Comparison: Treatment

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>		
<u>>140/90mmHg</u>		
<u>130-139/80-89mmHg</u>	<u>Lifestyle measures, maybe medication</u>	<u>Lifestyle measures</u>
<u>120-129/<80mmHg</u>	<u>Close monitoring</u>	<u>Close monitoring</u>
<u><120/80mmHg</u>	<u>Screening</u>	<u>Screening</u>

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
Gaseem et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinician Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. Ann Intern Med. 2017;166(6):430-437.

Guideline Comparison: Treatment

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>		
<u>>140/90mmHg</u>	<u>Medication</u>	<u>Lifestyle measures, maybe medication</u>
<u>130-139/80-89mmHg</u>	<u>Lifestyle measures, maybe medication</u>	<u>Lifestyle measures</u>
<u>120-129/<80mmHg</u>	<u>Close monitoring</u>	<u>Close monitoring</u>
<u><120/80mmHg</u>	<u>Screening</u>	<u>Screening</u>

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
Gaseem et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinician Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. Ann Intern Med. 2017;166(6):430-437.

Guideline Comparison: Treatment

	<u>AHA/ACC</u>	<u>ACP/AAFP</u>
<u>>150/90mmHg</u>	<u>Medication</u>	<u>Medication</u>
<u>>140/90mmHg</u>	<u>Medication</u>	<u>Lifestyle measures, maybe medication</u>
<u>130-139/80-89mmHg</u>	<u>Lifestyle measures, maybe medication</u>	<u>Lifestyle measures</u>
<u>120-129/<80mmHg</u>	<u>Close monitoring</u>	<u>Close monitoring</u>
<u><120/80mmHg</u>	<u>Screening</u>	<u>Screening</u>

Whelton PK et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. Hypertension 2017.
 Caseem et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinical Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. Ann Intern Med. 2017;166(6):439-437.

Conclusion

- Hypertension is prevalent
- Consequences are severe
- Proper diagnosis is key
- Intervention is effective
- Guidelines are a guide

Case #1

- 45 year-old African American man
- No medical problems
- Father and mother both had hypertension
- BMI = 25
- BP = 150/90

Case #2

- 50 year-old woman
- Just moved to your community
- Systolic heart failure; EF = 35%
- Current medication: diltiazem
- BP = 148/84

Case #3

- 65 year-old man with diabetes
- Medications: metformin & glyburide
- BP 136/86
- Creatinine = 1.8 with GFR = 35
- Urinalysis = 450 mg/dl protein

Case #4

- 70 year-old woman with paroxysmal atrial fibrillation causing palpitations and dizziness
- Episodes occur 1-2 times per month and last 10-20 minutes
- BP = 154/92

Case #5

- 30 year-old woman
- Essential hypertension diagnosed at 25
- Treatment: hydrochlorothiazide + Lisinopril
- Now 6 weeks pregnant
- BP = 120/70

Case #6

- 70 year-old man
- Normal health
- BP = 166/78

Case #7

- 55 year-old man
- Reports home BP < 130/80
- BP on arrival to clinic = 144/96