

Goals of sedation:

- 1. Patient safety**
- 2. Patient comfort**

Conscious Sedation

- Minimal Sedation
(anxiolysis)**
- Moderate Sedation**
- Deep Sedation**
- Anesthesia**

Minimal Sedation (Anxiolysis)

- **Patients respond normally to commands**
- **Cognitive function and coordination may be impaired**
- **Ventilatory and cardiovascular functions are unaffected**

Moderate Sedation


- **Depressed consciousness**
- **Patients respond purposefully to verbal commands**
- **No interventions are required to maintain airway**
- **Spontaneous ventilation is adequate**
- **Cardiovascular function is usually maintained**

Deep Sedation

- **Depressed consciousness**
- **Patients cannot be easily aroused but will respond after repeated or painful stimuli**
- **Ventilatory function may be impaired**
- **May required airway assistance**
- **Spontaneous ventilation may be inadequate**
- **Cardiovascular function is usually maintained**

General anesthesia

- **Patients are not arousable even with painful stimuli**
- **Ventilatory function is often impaired**
- **Often require airway assistance**
- **May require mechanical ventilation**
- **Cardiovascular function may be impaired**



**The sedation plan
must be clearly
articulated among
all members of the
procedure team**

Pre-sedation history

- Cardiac conditions
- Pulmonary conditions
- Renal disease
- Hepatic disease
- Endocrine disorders
- Head trauma
- Prior surgical or airway issues
- Prior intubation
- Stridor
- Snoring
- Sleep apnea
- Previous reactions to sedative medications

STOP-BANG

- | | |
|--|---|
| S – Snore: have you been told you snore | B – BMI: is your BMI greater than 28 |
| T – Tired: are you tired during the day | A – Age: 50 or over |
| O – Obstruction: do you stop breathing at night | N – Neck: circumference greater than 17 inches |
| P – Pressure: do you have high blood pressure | G – Gender: male |

Yes to 3 or more = increased risk for sleep apnea

Other key elements of the history:

- **Current medications**
- **Allergies**
- **Pregnancy status**
- **Last oral intake**
- **Need for isolation for infections**
- **Alcohol, tobacco, and drug use**

Physical examination

- **Cardiac exam**
- **Pulmonary exam**
- **Ability to lay in the proper procedure position**
- **Airway assessment**

ASA Physical Status

P1 - normal healthy patient

P2 – mild systemic disease

P3 – severe systemic disease

**P4 – severe systemic disease that is a
constant threat to life**

P5 – moribund and likely to die

P6 – brain dead organ donor

When to consider anesthesia consult?

- **Significant co-morbid disease**
- **Significant sleep apnea**
- **History of airway problems during sedation**
- **History of adverse reaction to sedation**
- **High risk airway**
- **Chronic opioid or sedative use**

Coding and billing for sedation

- For the physician doing both the procedure and the sedation:
 - ✓ 99152: Initial 15 minutes of sedation services
 - ✓ 99153: Each subsequent 15 minutes of sedation services
- For the physician doing only the sedation:
 - ✓ 99156: Initial 15 minutes of sedation services
 - ✓ 99157: Each subsequent 15 minutes of sedation services

Airway Assessment

John S. Rogoski, DO, FASA
Assistant Professor
Clinical Anesthesiology
The Ohio State University Wexner Medical Center

Four Types of Difficulty

- **Difficult to bag/mask ventilate/oxygenate**
- **Difficult laryngoscopy**
- **Difficult intubation**
- **Difficult to perform cricothyroidotomy**

How Does the ASA Define the Difficult Airway?

- **Difficult mask ventilation**
 - **Impossible for an unassisted anesthesiologist to prevent or reverse signs of inadequate ventilation during positive pressure mask ventilation**

How Does the ASA Define the Difficult Airway?

- **Difficult rigid laryngoscopy**
 - It is not possible to visualize any portion of the vocal cords with conventional laryngoscopy
- **Difficult intubation**
 - proper insertion of an endotracheal tube requires more than 3 attempts or greater than 10 minutes

Causes of Difficulty

- **Anatomical**
 - Obesity
 - Short neck
 - Protruding teeth, long high arched palate
 - Receding mandible
 - Decreased distance between occiput and spinous process
 - Increased alveolar-mental distance

Causes of Difficulty


- **Acquired**
 - **Acute neck swelling:** trauma, infection, post-operative bleeding
 - **Restricted jaw opening:** Trismus, fibrosis, rheumatoid arthritis, mandibular fracture
 - **Restricted neck movement:** osteoarthritis, scarring, C-spine tumor, ankylosing spondylitis

Predicting Difficult Bag & Mask Ventilation

- **B - bearded**
- **O - obese /obstetric**
- **N - no teeth**
- **E - elderly**
- **S - snores/sleep apnea**

Predicting Difficult Intubation Mallampati Classification

- **Class 1:** view of the entire posterior oropharynx to the bases of the tonsillar pillars
- **Class 4 :** no view of the posterior oropharynx or uvula



Korean J Pediatr. 2010 October; 53(10): 863–871.
Published online 2010 October 31

Predicting Difficult Intubation 3 - 3- 2 Rule

- **3 finger mouth opening**
- **3 fingers mentum to hyoid distance**
- **2 fingers hyoid to thyroid**

Predicting Difficult Intubation

- **Review medical record, history**
- **Assess**
 - **teeth especially protruding incisors**
 - **patent nares**
 - **open mouth & extend tongue (mallampati)**
 - **protrude mandible**
 - **thyromental distance, submental space**
 - **neck - short, thick ?, overall mobility & sniffing position**
 - **body habitus**


Video of Airway Examination



Airway Management

Supplemental Oxygen

- Nasal cannula
- Simple mask
- Non-rebreather mask



Creative Commons Attribution-ShareAlike 3.0 Unported

Airway Support

- Jaw thrust
- Nasal airways
- Oral airways



Bag / Mask Ventilation

- Technique dependent
- Mask seal essential
- 2 are better than 1
- Incorporate jaw thrust
- Nasal / Oral airways
- Assist spontaneous ventilation



Creative Commons
Attribution-ShareAlike 3.0
Unported



Video of Airway Maneuvers



Before the procedure

- **There must be signed written consent for:**
 - **The procedure**
 - **The sedation**
- **If 2 procedures are planned, get consent for both before giving sedation**
- **A “time-out” must be performed**

Q 5 minutes during the procedure:

- **Level of consciousness**
- **Blood pressure**
- **Oxygen saturation**
- **Respiratory rate**
- **Cardiac rhythm (only required in patients with known heart disease)**

Monitoring every 15 minutes until:

- **Patient is awake, alert, and oriented**
- **Recovered protective reflexes**
- **Vital signs returned to normal**
- **Oxygen saturation > 95% or at baseline**

Post-procedure transport:

- **Accompanying personnel trained in sedation monitoring**
- **Pulse oximeter**
- **Supplemental oxygen**
- **Ventilation equipment**
- **Nasal and/or oral airways**
- **Emergency drug supplies**
- **Cardiac monitor (in patients with heart disease)**

Post-procedure discharge:

- **Instruction sheet**
 - **No driving**
 - **No alcohol or sedatives**
 - **No operating machinery**
 - **Phone number for questions**
- **A responsible adult to accompany (taxi do not count!)**

Pharmacology of Sedatives & Reversal Agents

Mary Beth Shirk, PharmD, RPh
Clinical Associate Professor
The Ohio State University College of Pharmacy
Specialty Practice Pharmacist, Emergency Medicine
The Ohio State University Wexner Medical Center

Agents for Procedural Sedation

- Opioids
- Benzodiazepines
- Etomidate
- Ketamine
- Methohexital
- Propofol
- Dexmedetomidine

Opioids

- Class II Controlled Substances
- Mu receptor agonists
 - ✓ Fentanyl
 - ✓ Hydromorphone
 - ✓ Morphine
 - ✓ Meperidine
- Hepatic metabolism with varying $t_{1/2}$

Opioids

Adverse Effects

- Respiratory depression
- Hypotension
- Miosis
- Decreased GI motility
- Urinary retention

Opioids

Estimated IV Potency

- Fentanyl 75 - 100 micrograms
- Hydromorphone 1.5 mg
- Meperidine 75 mg
- Morphine 10 mg

Fentanyl

- Phenylpiperidine opioid agonist
- Preferred opioid for procedural sedation
- Precautions
 - ✓ Skeletal muscle and chest wall rigidity
 - Dose and administration rate related
 - Reversible with naloxone
 - ✓ Bradycardia
- Black box warning with CYP3A4 inhibitors

Benzodiazepines

- Class IV Controlled Substances
- GABA and Benzodiazepine agonists
 - ✓ Midazolam
 - ✓ Lorazepam
 - ✓ Diazepam
- Hepatic metabolism with varying $t_{1/2}$

Benzodiazepines

Adverse Effects

- Respiratory depression
- Hypotension
- Paradoxical reactions
- Nausea/vomiting
- Hiccoughs

Benzodiazepines

Estimated Potency

- Diazepam 5 mg
- Lorazepam 1 mg
- Midazolam 2 mg

Midazolam

- Preferred BZD for procedural sedation
- CYP3A4 substrate
- Elimination $t_{1/2}$ prolonged
 - ✓ CHF
 - ✓ Renal function impairment
 - ✓ Hepatic function impairment
 - ✓ Obesity
 - ✓ Elderly

Etomidate

- Not currently controlled substance
- Nonbarbiturate benzylimidazole hypnotic
- 0.1 – 0.3 mg / kg IVP *over 30-60 seconds*

Etomidate

- Inhibits 11- β hydroxylase
- Blocks cortisol production
- Myoclonus (up to 33%)
- Injection site pain (30-80%)
 - ✓ Propylene glycol
- Minimal effect on hemodynamics
- Decreases ICP and seizure threshold

Ketamine

- Class III Controlled Substance
- NMDA receptor antagonist and PCP derivative
- Analgesic properties appealing
- IM or IV administration
- 0.5 – 2 mg/kg IVP *over at least 60 seconds*

Ketamine

- Respiratory drive maintained
- Three concentrations available
 - ✓ 10 mg/mL
 - ✓ 50 mg/mL
 - ✓ 100 mg/mL (dilute if administered IV)

Ketamine

Emergence reaction (12 - 50%)

- ✓ Severity varies
- ✓ Less common in < 15 yrs and > 65 yrs
- ✓ Less frequent with IM administration
- ✓ Minimize verbal, tactile, visual stimulation during recover
- ✓ ?pretreat with BZD or butyrophenone

Ketamine

- Emergence reaction (12- 50%)
- Hypersalivation ? pretreat?
- Nystagmus
- Increases ICP/IOP
- Minimal affect on BP/HR or increase
- Increased skeletal muscle tone

Methohexital

- Class IV controlled substance
- Ultrashort acting IV barbiturate anesthetic
- pH of 1% solution is 10-11
- Contraindicated in porphyria
- Hypotension
- Respiratory depression
- Dose 0.25 – 1 mg/kg *at* <10mg/5 seconds
- 500 mg vials!

Propofol

- Currently not controlled substance
- Patient can transition in unpredictable fashion to deeper level of sedation
- At OSUWMC physician must be credentialed for deep sedation
- Cardiovascular depressant – hypotension!

Propofol

- Contraindicated if
 - ✓ egg allergy (?)
 - ✓ soy intolerance (?)
 - ✓ peanut allergy (?)
- 0.5 - 1 mg/kg IV over 2-3 *min* once then 0.5 mg/kg every 3-5 *min* if needed

Dexmedetomidine

- “relatively selective” α_2 adrenergic agonist
- FDA approval in 2008
 - ✓ Sedation of nonintubated patients prior to and/or during surgical and other procedures
- Limited published experience for procedural sedation (ablation, hysteroscopy, etc)

Dexmedetomidine

- Hypotension 54% vs 30% (Placebo)
 - ✓ SBP<80 or DBP <50 or ↓ >30% from baseline
 - ✓ 72% in ≥ 65yo patients (n=131)
- Bradycardia/sinus arrest 14% vs 4% (Placebo)
 - ✓ <40BPM or ↓ >30% from baseline

Approaches Being Explored

- Alternative routes of administration
 - ✓ Intranasal
 - ✓ Nebulized
- Alternative combinations of medications
 - ✓ Ketamine + Propofol
 - ✓ Ketamine + Dexmedetomidine

	Onset (Min)	Peak (Min)	Duration (Min)	Elimination
Fentanyl	Immed	Immed	30-60	Hepatic
Midazolam	1-2	2-2.5	30	Hepatic + (Renal)
Etomidate	<1	1	3-5	Hepatic
Ketamine	1	1	15-20	Hepatic Active Metabolite
Methohexital	Immed	Immed	10-20	Hepatic
Propofol	½	1	3-10	Hepatic
Dexmedetomidine			4 hours	Hepatic

	Amnestic	Analgesic	Anxiolytic
Benzodiazepines	+	-	+
Opioids	-	+	-/+
Etomidate	+	-	+
Ketamine	+	+	Dissociative properties
Methohexital	-	-	+
Propofol	+/-	-	+
Dexmedetomidine	+	+	+

Recommended Agents at OSUWMC

- Midazolam ± fentanyl agents of choice
- Propofol limited to physicians credentialed in deep sedation
- Meperidine not for routine use
- Alternative agents used by physician experienced in their use

Dose

- No universally safe & effective dose
- Variable dose requirements
 - ✓ Age (especially >65 yrs)
 - ✓ Weight
 - ✓ Medical condition
 - ✓ Medication history
 - ✓ Previous requirements during procedures
 - ✓ Goal depth of sedation

Dose

- Combination agents have added risks/benefits
- TITRATE
 - ✓ Small incremental doses
 - ✓ ***Sufficient time must elapse*** between doses to evaluate effect of previous dose
 - ✓ Time between doses longer for nonintravenous routes

Fentanyl: Typical Initial Regimen*

- 25-100 micrograms SLOW IVP
- IVP over *at 1 - 2 minutes*
- Dilute to permit slower administration
- *Additional doses in 2 minutes if needed*
- Administer prior to midazolam if using combination regimen

*Dose is highly variable

Midazolam: Typical Initial Regimen*

- 0.2 – 2.5 mg IVP
- IVP over *at least 2 minutes*
- Dilute to permit slower administration
- *Additional dose(s) in 3 minutes if needed*
- Administer after opioid if using combination regimen

*Dose is highly variable

JCAHO & Medication Administration During Procedures

- Sterile technique!
- Proper product labeling
 - ✓ Label: drug name, strength, and amount
 - ✓ Single individual process and immediate administration = no label
 - ✓ Two individual process = product verification with vial and label

JCAHO & Medication Administration During Procedures

- Document waste of Controlled Substances
- Complete charting
 - ✓ Medication
 - ✓ Dose
 - ✓ Route
 - ✓ Time of administration
 - ✓ Who administers

Reversal Agents

- Used to reverse sedatives or treat overdose
- Half lives can be shorter than sedative
- Can precipitate withdrawal symptoms
- May not completely reverse all complications of sedatives

Flumazenil

- Onset of action 1-2 minutes
- Half life 41-79 minutes
- Flumazenil use requires 90 min monitored recovery time
- Hepatic clearance

Flumazenil

- Adverse Effects
 - ✓ Seizures
 - ✓ Panic attacks and emotional lability
 - ✓ Withdrawal symptoms
 - ✓ Dizziness
- Reversal of Procedural Sedation
 - ✓ 0.2mg IVP q 1 min prn to MAX of 1mg
 - ✓ Repeat every 20 min as needed

Naloxone

- Opiate receptor antagonist
- Onset of action 2-3 minutes
- Half life 30-81 minutes
- Naloxone use requires 90 min monitored recovery time
- Duration of effect varies (45min – 4 hrs)
- Hepatic clearance

Naloxone

- Dosing
 - ✓ 0.1 – 0.2 mg IVP every 1-2 minutes
 - ✓ Doses up to 2 mg may be required
 - ✓ May need to redose if naloxone wears off before opiate
- Adverse Effects
 - ✓ Opiate withdrawal
 - ✓ Pulmonary edema
 - ✓ Acute hypertension and dysrhythmias
 - ✓ Seizures

Moderate and Deep Sedation

Deep sedation

- **Emergency medicine**
- **Pulmonary medicine**
- **Critical care**
- **Oral maxillary facial surgery**
- **Or demonstrated advanced airway expertise and intubation skill**