



# Systemic Lupus Erythematosus: Diagnosis and Management

**Judith Lin, MD**

*Assistant Professor of Medicine*

*Division of Rheumatology*

*Department of Internal Medicine*

*The Ohio State University Wexner Medical Center*

**MedNet21**  
Center for Continuing Medical Education

 **THE OHIO STATE UNIVERSITY**  
WEXNER MEDICAL CENTER

## Disclosures

- None

## Objectives

1. Identify clinical features and common manifestations of SLE
2. Identify immunologic findings of SLE
3. Recognize common SLE treatments and associated side effects
4. Recognize complications that may be seen with SLE and the importance of health maintenance management

## What is SLE?

- **Systemic autoimmune** disease characterized by heterogenous **multisystem** involvement and production of **autoantibodies**
  - Driven by **loss of immune tolerance** and abnormal innate and adaptive immune function
  - Immune complex mediated reactions and tissue destruction
  - Variable clinical presentation and clinical course

## Risk factors for SLE

- Women of childbearing age
- More in African American, Hispanic, other ethnic minorities
- Genetics
  - Polygenic
  - Early complement deficiencies
  - Family history
- Environment
  - Infections, **smoking**, UV exposure, drugs, stress
- Genetics + environment → Immune dysregulation

## Diagnosis vs Classification

### Diagnosis

- based on **clinical presentation** combined with serologic findings
  - +ANA is not enough
- No diagnostic criteria
- Diagnosis made by experienced physician/rheumatologist

### Several Classification criteria

- ACR/EULAR, SLICC
- For categorizing patients for research purposes
- Not intended as diagnostic criteria
  - Can be used as guide to organize thoughts

Exclude alternative diagnosis

## 1997 ACR Classification criteria

4 or more criteria, excluding other causes

Criteria	Definition
Malar rash	Fixed erythema over malar eminences sparing nasolabial fold
Discoid rash	Erythematous raised patches with adherent keratotic scale and follicular plugging, often with atrophic scars
Photosensitivity	Rash from unusual reaction to sunlight
Oral ulcers	Oral or nasopharyngeal ulcers, usually painless, observed by physician
Arthritis	Nonerosive, 2 or more peripheral joints with tenderness, swelling or effusion
Pleuritis or pericarditis	Convincing history or objective evidence
Renal disorder	Persistent proteinuria, >0.5g/24hr or >3+ on dipstick, cellular casts
Neurologic disorder	Seizures or psychosis in the absence of offending drugs or metabolic derangements
Hematologic disorder	Hemolytic anemia, leukopenia, lymphopenia, or thrombocytopenia
Immunologic disorder	Anti-dsDNA, Sm, or antiphospholipid antibodies
Positive ANA	Abnormal titer at any point in time, in absence of drugs known to be associated with drug induced lupus

Arth Rheum 1997

## 2012 SLICC Classification criteria

Clinical Criteria	Immunologic criteria	
Acute cutaneous lupus: malar rash, SCLE, others	ANA above lab reference range	≥4 criteria:
Chronic cutaneous lupus • Discoid, panniculitis, lupus tumidus, chillblains	Anti-dsDNA above lab reference range, except ELISA: twice above lab reference range	
Oral ulcers: palate	Anti-Smith	• 1 clinical <u>and</u>
Nonscarring alopecia		• 1
Synovitis involving 2 or more joints	Low complement (C3, C4, CH50)	immunologic
Serositis: Pleuritis, pericarditis	Direct Coombs test in the absence of hemolytic anemia	• exclude other causes
Renal disorder • UPCR or 24hr urine protein ≥500mg/24hr • RBC casts		Lupus nephritis can be made by biopsy and +ANA alone
Neurologic • Seizures, psychosis • Mononeuritis multiplex • Myelitis • Peripheral or cranial neuropathy • Acute confusional state	Antiphospholipid antibody: any of the following: • Lupus anticoagulant • False positive RPR • Medium or high titer anticardiolipin (IgG, IgM, or IgA) • Anti-B2 glycoprotein I (IgG, IgM, or IgA)	
Hemolytic anemia		
Leukopenia (<4000/mm <sup>3</sup> ) or lymphopenia (<1000/mm <sup>3</sup> )		
Thrombocytopenia (<100,000/mm <sup>3</sup> )		

Arth Rheum 2012

## 2019 EULAR/ACR SLE Classification Criteria

**Entry criteria: ANA  $\geq$ 1:80**

**Additive criteria: at least 1 clinical and  $\geq$ 10 points**

- Only the highest weighted criteria is scored within each domain
- Criteria does not need to be simultaneous

Exclude alternative causes

Aringer M. Arth Rheum 2019


Clinical domains and criteria	Weight
<b>Constitutional</b>	
Fever	2
<b>Hematologic</b>	
Leukopenia	3
Thrombocytopenia	4
Autoimmune hemolysis	4
<b>Neuropsychiatric</b>	
Delirium	2
Psychosis	3
Seizure	5
<b>Mucocutaneous</b>	
Non-scarring alopecia	2
Oral ulcers	2
Subacute cutaneous OR discoid lupus	4
Acute cutaneous lupus	6
<b>Musculoskeletal</b>	
Joint involvement	6

Clinical domains and criteria	Weight
<b>Serosal</b>	
Pleural or pericardial effusion	5
Acute pericarditis	6
<b>Renal</b>	
Proteinuria $>$ 0.5g/24hr	4
Renal biopsy class II or V LN	8
Renal biopsy Class III or IV LN	10

Immunology domains and criteria	Weight
<b>Antiphospholipid antibodies</b>	
Anti-cardiolipin antibodies OR Anti-B2GPI antibodies OR Lupus anticoagulant	2
<b>Complement</b>	
Low C3 OR low C4	3
Low C3 AND low C4	4
<b>SLE-specific antibodies</b>	
Anti-dsDNA or Smith antibody	6

## Antinuclear antibody (ANA)

- Antibodies against proteins or nucleic acids in nucleus
- Found in  $>$ 95% of SLE but only 57% specific
- Detection assays
  - Indirect immunofluorescence (IIF)
    - Gold standard
    - Titer
    - Staining pattern may guide clinical thinking
    - Time consuming, labor intensive, may have false positive
  - ELISA
    - Antibodies to different nuclear antigens
    - Faster, detect specific antibodies
    - High sensitivity but less specific



By Simon Caulton - Own work, CC BY-SA 3.0, <https://commons.wikimedia.org/w/index.php?curid=20521932>

ANA more likely to have clinical significance with titers  $\geq$ 1:80

ANA is sensitive but not specific for SLE, higher titer more likely to be associated with autoimmune disease


## ANA titer and prevalence

- ANA common in general population
  - 25-30% have 1:40 titer
  - 10-15% have 1:80 titer
  - 5% have 1:160 titer or higher

Solomon DH et al. Arth&Rheum 2002

ANA is common in healthy subjects

## ANA Prevalence increases with age



Satoh M, Chan EK, Ho LA, et al. Prevalence and sociodemographic correlates of antinuclear antibodies in the United States. *Arthritis Rheum.* 2012;64(7):2319-2327. Copyright obtained from publisher.

## ANA is common and nonspecific

ANA is seen in various conditions

- Can be triggered by
  - Infections
  - Smoking
  - Silica, other chemicals and pollutants
  - Medications:
    - Hydralazine
    - Procainamide
    - Isoniazid
    - Minocycline
    - TNF alpha inhibitors
- Can be seen in other conditions:
  - Other autoimmune disease
    - Other systemic autoimmune rheumatic disease
    - Hashimoto's thyroiditis
    - Multiple sclerosis
    - Psoriasis
    - Autoimmune hepatitis
    - Idiopathic thrombocytopenic purpura
  - Atopic diseases
  - Infections
  - Malignancies
  - Liver disease
  - Family history of autoimmune disease

Soloman DH et al. Arth Care & Res 2002

Recognize common autoantibodies in SLE

## ANA subsets/Extractable nuclear antigen antibodies (ENAs)

Antibody	Frequency	ANA pattern	Clinical associations
Smith	30%	Speckled	High specificity, low sensitivity. More in African Americans. More organ damage.
dsDNA	70%	Homogenous	Can fluctuate with disease activity. Gold standard Crithidia assay is very specific, but common ELISA assay not very specific
SSA	30%	Speckled	Sicca, photosensitivity. Seen with Sjogren's, SCLE, NNL, CHB
SSB	20%	Speckled	SCLE, Sjogren's, NNL, CHB
U1RNP	25-40%	Speckled	MCTD, Raynaud, ILD, pulmonary hypertension
Histone	50%	Homogenous	SLE and DIL (75%)

NNL: neonatal lupus, CHB: congenital heart block

### Antiphospholipid antibodies (aPL)

- Lupus anticoagulant
- B2glycoprotein I IgG, IgM
- Cardiolipin IgG, IgM

### Complements

- C3
- C4
- CH50
- C1q

*Antibody specificity and sensitivity limited by commercial assays*

### Choosing wisely campaign:

Avoid ordering ANA sub-serologies if ANA negative and low clinical suspicion of immune-mediated disease

Exceptions: Jo1 and SSA

**Antibodies alone are not sufficient to make diagnosis**

Consider other lupus-related diseases

## Other forms of lupus and lupus-related disorders

- Cutaneous lupus
- Neonatal lupus
- Mixed connective tissue disease
- Antiphospholipid syndrome
- Sjogren's syndrome
- Undifferentiated connective tissue
- Overlap syndrome

### Drug-induced lupus

- Hydralazine
- Propylthiouracil
- Sulfonamides
- Lithium
- Anticonvulsants
- Quinidine
- Diltiazem
- Beta blockers
- Interferon gamma
- TNF inhibitors

## SLE Management

### SLE treatment goals

- Control disease activity
  - Goal of remission or low disease activity
- Minimize complications from disease and treatment
- Improve quality of life

### Preventative measures

- **Smoking** cessation
- **Photoprotection**
- Avoid medications that may trigger lupus if possible

### Treat reversible causes of symptoms

- Physical and lifestyle measures
  - Address fatigue, sleep, exercise
  - Provide emotional and psychosocial support
  - Assess and treat fibromyalgia

### Treat associated comorbidities

- Other autoimmune disease: T1DM, Hashimoto's

### Health maintenance

- **Cardiovascular** health assessment
  - Assess and treat reversible risk factors given increased risk of **CVD**
- **Bone** health assessment
  - Increased risk of **osteoporosis, avascular necrosis** due to SLE, sun avoidance, steroid use
- Age-appropriate **cancer screening**
  - Increased risk of **malignancy** in SLE
- **Immunizations**
- **Contraception** counseling
- **Pregnancy** planning

All SLE patients require multi-disciplinary care with PCP, rheumatology and other specialists to optimize management and outcomes

## Systemic corticosteroids

- For rapid control of inflammatory activity
- Usually given as taper
- Pulse dose
  - IVMP for severe organ threatening disease
- High dose
  - For severe disease such as serositis, nephritis, hemolytic anemia
  - Prednisone 20mg or higher
- Moderate dose
  - For moderate disease such as arthritis
  - Prednisone 7.5-20mg
- Low dose:
  - Usually used as slow taper or maintenance
  - prednisone 7.5mg or lower

Steroids can be given for SLE flares but limit use as it is associated with significant side effects

### Side effects:

- Osteoporosis, avascular necrosis, bone fractures
- Weight gain, Cushingoid features
- Hyperglycemia, diabetes
- Fluid retention, hypertension
- Arrhythmia
- Cataracts, glaucoma
- Gastritis, PUD
- Mood disorder, psychosis
- Muscle weakness
- Adrenal insufficiency
- Skin thinning, ecchymosis, striae



## Anti-malarials

Antimalarials reduce flares and improves outcomes in SLE

### Hydroxychloroquine, Chloroquine, Quinacrine

- For active, non-organ threatening SLE
  - Rash, arthritis, alopecia
- Many benefits in SLE:
  - Reduce risk of flares in SLE
  - Prevent progression of disease
  - Reduce thrombotic and cardiovascular complications
  - Improve glucose and lipid profiles
- Slow onset of action
  - Weeks to months to see effect
- Hydroxychloroquine dose
  - Up to 5mg/kg/day (max 400mg/day)
  - Dose reduce for renal insufficiency

### Side effects

- **Retinal toxicity:**
  - Risk increases with time
  - Irreversible
  - **Need retinal exam yearly**
- Drug rash
- Blue-gray discoloration of skin
- GI upset
- Myopathy
- Cardiomyopathy
- Arrhythmia
- CNS disturbance (dizziness, headache, insomnia, psychosis)
- **Caution in G6PD deficiency**
- **Safe in pregnancy**

## Systemic immunosuppressive therapy

- Cytotoxic therapy
  - **Azathioprine**
  - **Mycophenolate**
  - (Methotrexate)
  - (Leflunomide)
  - **Cyclophosphamide**
- Calcineurin inhibitors
  - (Cyclosporin)
  - (Tacrolimus)
  - Voclosporin
- Biologic therapy
  - **Belimumab**
  - (Rituximab)
  - (Others to come)

## Case #1

- 40 yo Hispanic F w/ h/o HTN, anemia
- Joint pain and swelling in hands with morning stiffness
- Facial rash and body rashes
- Alopecia, oral ulcers
- No smoking
- Fam hx: No autoimmune disease



Jaccoud's arthropathy

## Case #1: Workup and diagnosis

- +ANA 1:320 speckled
- +Sm, +SSA
- +ribosomal P, +chromatin
- +dsDNA 48
- +RF 27, -CCP
- Cq1 7 (L) → early complement deficiency increases risk of SLE
- C3 39, C4 <8 (L)
- CBC with ACD otherwise normal
- ESR, CRP normal
- UA, UPCR normal
- Skin biopsy: interface dermatitis

### Diagnosis:

**SLE** (+ANA, +Sm, +SSA, +ribosomal P, +chromatin, +dsDNA, +RF) with **hypocomplementemia, anemia, Jaccoud's arthropathy, oral ulcers, alopecia, acute cutaneous lupus**

# Cutaneous lupus

## Acute cutaneous lupus

Localized: Malar rash

- Distinguish from:
  - Rosacea
  - Seborrhea dermatitis
  - Dermatomyositis

Generalized

## Subacute cutaneous lupus erythematosus (SCLE)

- Types
  - Annular
  - Papulosquamous
- Photosensitive
- 50% have SLE
- 70% with +ANA, +SSA, 30% SSB



## Different forms of cutaneous LE

### Chronic cutaneous lupus

- Discoid lupus
- Lupus profundus (panniculitis)
- Chilblain
- Tumid lupus

### Other cutaneous manifestations

- Raynaud
- Vasculitis
- Livedo reticularis
- Urticaria
- Others

# Alopecia

## Non-scarring alopecia

- Focal or diffuse
- Differential:
  - Traction alopecia
  - Female pattern hair loss
    - crown, frontal, hereditary
  - Telogen effluvium
  - Iron deficiency
  - Hypothyroidism



## Recognize alopecia in SLE

### Scarring alopecia

- Inflammatory, infiltrative conditions
- More focal than diffuse
- Discoid lupus



## Case #1: Treatment

Belimumab used as add-on therapy for SLE

- Topical therapy for cutaneous lupus
  - Topical corticosteroids
  - Topical calcineurin inhibitors
- Immunomodulator
  - Hydroxychloroquine
- Clinical course:
  - Inadequate control on plaquenil
  - Intolerant to azathioprine, mycophenolate
  - Started on belimumab
    - Complicated by infection

### Belimumab

- Monoclonal ab against BLYS
- SQ and IV
- For active seropositive SLE
  - Best for +dsDNA, low complements, skin, MSK manifestations
- Adverse effects
  - Infection
  - Injection site/infusion reaction
  - Diarrhea, nausea
  - Headache
  - Psych: depression, suicidal ideation
  - Cytopenias
  - PML

## Infection evaluation, management, prevention

	Infection	SLE flare
WBC	↑	↓
ESR	↑	↑
CRP	↑	--/↑
C3, C4, CH50	--/↑	↓
dsDNA	--	↑

### ***SLE flare can be triggered by infections***

#### Management during infection



- Hold immunosuppressive therapies until infection resolves
- Ok to continue:
  - Hydroxychloroquine
  - Prednisone if chronic

### Vaccinations

- Recommended vaccines on immunosuppressive therapy:
  - Yearly influenza
  - Prevnar13
  - Pneumovax23
  - HAV and HBV
  - HPV
  - Shingrix
- Avoid live vaccines
- Vaccines may be more effective when given before starting immunosuppressive therapy

## Case # 2

- 40 yo AAF with h/o cervical lymphadenopathy (biopsy negative for malignancy) who presented with SOB and leg swelling
- SOB worse with lying down
- Smoker
- Exam
  - BP 115/80, HR 101, RR 20, 96% RA
  - periorbital edema, diffuse anasarca, ascites
  - Distant heart sounds
  - Rash on trunk, extremities, hands and feet



EKG: sinus tach, low voltage

## Chest pain, SOB in SLE

Differential for cardiac and pulmonary manifestations in SLE

### Cardiac manifestations


- Pericarditis
- Myocarditis
- Libman-Sacks endocarditis
- Coronary arteritis
- Arrhythmia
- CAD/MI
  - Accelerated atherosclerosis
    - 2+ fold increase risk of CAD, CVA, PAD

### Pulmonary manifestations

- Pleuritis
- Parenchymal lung disease
  - Pneumonitis
  - Diffuse alveolar hemorrhage
  - Interstitial lung disease
- Pulmonary vascular disease
  - Pulmonary hypertension
  - Pulmonary embolism
- Shrinking lung syndrome

Sent to hospital

- WBC 3.02 (L), abs lymph 0.4 (L) → concern for SLE flare
- Hb 8.3 (L), MCV 85
- Plt 175 → less likely hemolytic
- Troponin normal → less likely MI, myocarditis
- Cr 0.84 (baseline; BMP normal)
- UA trace proteins, no casts → less likely GN
- LFT normal except Albumin <1.5 (L)  
→ Protein loss leading to anasarca
- Chest XR: cardiomegaly, left pleural effusion  
→ Serositis from SLE
- TTE: large, circumferential pericardial effusion with early signs of tamponade → serositis from SLE (No endocarditis)



CT a/p: bowel wall edema  
→ Concern for protein-losing enteropathy

## Case #2

Serologies

- ANA >1:1280 speckled
- +Sm, +RNP, +SSA, +SSB
- +dsDNA 70 (H)
- Low C3 39, C4<8
- IgG 384 (L), IgA 59 (L), IgM normal.
- ESR 91 (H), CRP normal
- UPCR 0.484
- Stool alpha1antitrypsin 310 (H)

Serologies suggestive of SLE

- Pericardiocentesis and paracentesis
  - Exclude infection, malignancy
- Colonoscopy excluded alternative causes

**Diagnosis:**

**SLE** (+ANA, +Sm, +RNP, +SSA, +SSB, +dsDNA, low complements, leukopenia, lymphopenia, anemia) with **acute cutaneous lupus, pericarditis, pleuritis and protein-losing enteropathy**

## Case #2: Management

Cyclophosphamide effective induction therapy for severe SLE but has significant toxicities

### Treatment:

- **IVMP** for rapid control
  - Followed by PO taper
- **Hydroxychloroquine**
- **IVIG**
  - for hypogammaglobulinemia
- **Cyclophosphamide**
  - For induction therapy for severe organ threatening disease
  - IV vs PO


### Cyclophosphamide

- Used as Induction therapy for severe organ damage
- Toxicity increases with cumulative dose
- Transition to alternative agent for maintenance
- Adverse effects:
  - Hemorrhagic cystitis, transitional cell carcinoma
  - Cytopenia: **Leukopenia/neutropenia**
    - Monitored closely and adjust dose
  - GI upset, mucositis, stomatitis
  - Alopecia
  - **Gonadal failure, teratogenic**
    - Fertility discussion with obgyn prior to initiation

## Case #3

40 yo Caucasian F w/ h/o Factor V Leiden and h/o DVT who presented with blurry vision, found to have retinal hemorrhages by ophthalmology and admitted to hospital for hypertensive emergency

Chest XR: Small bilateral pleural effusions, mild bibasilar airspace disease, marked cardiomegaly



- BP 215/126
- WBC 2.42 (ALC 0.6), Hb 6.8, plt 112
- Retic 2.79%, haptoglobin <30, LDH 265
- Peripheral smear +schistocytes
  - hemolytic anemia
- ADAMTS13 activity 51% (normal >68%)
- UA: RBC, proteins. No casts.
- Cr 2.4 (baseline 1)
  - TTP/HUS, TMA, GN
- LFT normal except albumin 2

Pancytopenia, serositis, hemolytic anemia, AKI, multisystem organ involvement suspicious for SLE

- ANA 1:320 speckled
- +SSA, +SSB, +dsDNA 380
- APS negative, negative DAT
- C3 38, C4 <8 (L)
- Normal ESR, CRP
- UPCR 4.2g/24hr → nephrotic syndrome, LN

## Hematologic manifestations in SLE

- Leukopenia/lymphopenia
- Anemia
  - Hemolytic anemia
  - Anemia of chronic disease
- Thrombocytopenia
  - ITP, TTP
- Thrombotic microangiopathy
- Lymphadenopathy
- Splenomegaly
- Thromboembolism → check for APS



### Cytopenias:

- disease vs medications vs alternative causes

### Hemolytic anemia

- Autoimmune
- TTP/HUS
- DIC
- APS
- Other: valves, malignant hypertension, PNH

## Lupus nephritis

- Suspect in SLE with AKI, proteinuria, hematuria, active urinary sediment, hypertension
- 50% of SLE, high morbidity and mortality
- More common and more severe in Black and Hispanic

- **Role of kidney biopsy**
  - Establish diagnosis
  - Evaluate for other causes
  - Results determine treatment
- **Indication for biopsy**
  - Increase Cr without clear cause
  - Proteinuria >0.5g/24hr with active urinary sediment

### ISN/RPS 2003 Classification of LN

Class	Histologic classification
Class I	Minimal mesangial LN
Class II	Mesangial proliferative LN
Class III	Focal LN (<50% of glomeruli)
Class IV	Diffuse (>50% glomeruli) Diffuse segmental or global
Class V	Membranous LN
Class VI	Advanced sclerosing LN (>90% sclerosed glomeruli globally)

### **Proliferative lupus nephritis (class III and IV)**

#### Induction

- High dose steroids
- Cyclophosphamide
- Mycophenolate\*

#### Maintenance

- Mycophenolate
- Azathioprine
- Calcineurin inhibitors

\* African Americans respond better to MMF than CYC induction for lupus nephritis

Hahn BH. Arthr Care Res 2013



## Case #3: Diagnosis and management

### Renal biopsy:

- Diffuse proliferative lupus nephritis (Class IV)
- Active thrombotic microangiopathy
- Moderate interstitial fibrosis and tubular atrophy

### Diagnosis:

- **SLE** (+SSA, +SSB, +dsDNA, low complements), with pancytopenia, serositis
- **Class IV nephritis**
- **Thrombotic microangiopathy (aHUS)**

### Treatment:

- **IVMP** 1gx3 days, followed by PO prednisone
  - For rapid control of LN and hemolysis
  - Prophylaxis: Bactrim, PPI, calcium/vit D
  - Screen for hepatitis B/C, TB
- **Hydroxychloroquine**
- **Mycophenolate**
  - For LN after cell counts recover
- **Anticoagulation** with heparin transition to coumadin
  - For TMA
- **Eculizumab** for aHUS

## Antimetabolites

For inflammatory lung disease, lupus nephritis, other deep organ involvement

- First line therapy

For cutaneous lupus, joints

- Used after topical and anti-malarials

### Mycophenolate \*

- Inhibits purine synthesis
- PO 2-3g/day in BID dosing
- Common adverse effects
  - Infection
  - GI upset
  - Cytopenias
  - Elevated LFTs
  - PPI may reduce absorption
  - **Teratogenic**
  - OCP may be less effective

### Azathioprine \*

- Purine analog
- PO 2-2.5mg/kg/day
- Common adverse effects
  - Infection
  - GI upset
  - Cytopenia
  - Elevated LFTs
  - Headaches
- **Avoid in poor TPMT metabolizers**
- Safe in pregnancy

**\*Increase risk of malignancy**

### Recommend:

- **Age-appropriate cancer screening**
- **High vigilance**
- **HPV vaccine**

Mycophenolate and azathioprine are commonly used in SLE as first line therapy for deep organ involvement but may carry an increased risk of malignancy

## Case 4




- 31 yo Asian F w/ h/o SLE diagnosed in 2008 (+ANA >1:2560 speckled, +dsDNA, +Sm, +RNP, +RF, low C3/C4) with Raynaud, cutaneous ulcers and vasculitis with digital gangrene s/p amputation
- h/o non-adherence to medications and lost to follow up
- Presented with confusion and worsening cutaneous vasculitis

Exam: cushingoid, malar rash, distal vasculitic purpura, livedo reticularis, Raynaud, flat affect, hyper-reflexia in the lower extremities, up going toes bilaterally

## Case #4: Workup

- Admitted and found to have bilateral basal ganglia strokes



- WBC 2.5 (ALC 0.8) Hb 10.9, plt 209
- ESR 40, CRP 10.3 (H)
- dsDNA 35 (H)
- C3 59, C4 10 (L)
- Lupus anticoagulant positive
- B2GPI IgG 196, IgM 24
- Cardiolipin IgG 48, IgM normal
- UA, UPCR normal
- CSF with mildly elevated proteins and WBC, oligoclonal bands
- Ruled out infectious etiologies

## Neuropsychiatric lupus

Depression, anxiety	}	Neuropsychiatric evaluation
Psychosis		MRI of brain or spine
Cognitive dysfunction		CSF studies
Aseptic meningitis		EEG
Cerebritis/cerebral vasculitis		Autoantibodies:
CVA/TIA		• dsDNA, NMDA, NMO, ribosomal P, APS
Seizures		EMG/NCV
Demyelinating syndromes		
Transverse myelitis		
Neuromyelitis optica		
Peripheral neuropathy		NCV
Cranial nerve palsy		
Autonomic disorder		Autonomic testing
Headache		

## Case #4: diagnosis and management

### Diagnosis

**SLE** with cutaneous vasculitis and encephalopathy, ischemic CVA associated with **neuropsychiatric lupus** and secondary **antiphospholipid syndrome (APS)**

### Treatment

- IVMP followed by PO prednisone
- Cyclophosphamide
- Hydroxychloroquine
- Aspirin and anticoagulation
- Neurocognitive rehab
- Wound care
- Physical therapy
- *Close follow up to ensure adherence*

### Thromboembolic risk increased in SLE

- 2x risk of ischemic CVA in SLE
- +aPL increases risk of thrombosis
  - 40% SLE patients have aPL
- Evaluate and treat modifiable risk factors
  - Lifestyle changes
  - Hypertension
  - Hyperlipidemia
  - Smoking cessation
  - Avoid estrogen-containing contraceptives
- Treatment
  - Low dose aspirin
  - Warfarin preferred over DOACs
  - Hydroxychloroquine

SLE and aPL/APS increases risk of thromboembolism

## Summary

- Clinical manifestations of SLE
  - Cutaneous lupus
  - Cardiopulmonary manifestations
  - Lupus nephritis
  - Hematologic abnormalities
  - Neuropsychiatric lupus
  - Thromboembolism
    - Increased risk of CAD, CVA, PAD
    - +aPL/APS increases risk
- Immunologic findings
  - ANA and ENAs
  - Complements
  - Inflammatory markers
- Treatments
  - Antimalarials
  - Glucocorticoids
  - Mycophenolate, azathioprine, cyclophosphamide, belimumab
- Complications
  - Infections
  - Cardiovascular disease
  - Osteoporosis
  - Malignancy

## Take home points

- SLE can present with a wide range of clinical manifestation and diagnosis should be made by an experienced physician based on clinical presentation excluding alternative diagnosis and supported by serologic findings
  - Positive ANA is common and not sufficient to establish a diagnosis
- Treatment of SLE depends on areas being affected, disease activity and severity. Treatments may be associated with various toxicities that need to be monitored closely.
  - Hydroxychloroquine improves outcomes in SLE
- SLE patients have increased risk of infections (if on immunosuppressive therapy), cardiovascular disease, thromboembolism, renal disease, osteoporosis, and malignancy
  - Patient should be counseled on immunizations and infectious management, evaluated and treated for cardiovascular and thromboembolic risk factors, screened for lupus nephritis, evaluated for bone health, counseled on contraception with the use of teratogenic medications, and follow appropriate cancer screenings.

## References

- Okon LG, Werth VP. Cutaneous lupus erythematosus: diagnosis and treatment. *Best Pract Res Clin Rheumatol*. 2013;27(3):391-404. doi:10.1016/j.berh.2013.07.008
- Kuhn A, Bonsmann G, Anders HJ, Herzer P, Tenbrock K, Schneider M. The Diagnosis and Treatment of Systemic Lupus Erythematosus. *Dtsch Arztebl Int*. 2015;112(25):423-432. doi:10.3238/arztebl.2015.0423
- Hahn, B. H., McMahon, M. A., Wilkinson, A., Wallace, W. D., Daikh, D. I., Fitzgerald, J. D., Karpouzas, G. A., Merrill, J. T., Wallace, D. J., Yazdany, J., Ramsey-Goldman, R., Singh, K., Khalighi, M., Choi, S. I., Gogia, M., Kafaja, S., Kamgar, M., Lau, C., Martin, W. J., Parikh, S., ... American College of Rheumatology (2012). American College of Rheumatology guidelines for screening, treatment, and management of lupus nephritis. *Arthritis care & research*, 64(6), 797–808. <https://doi.org/10.1002/acr.21664>

## ANA references

- Abeles AM, Abeles M. The clinical utility of a positive antinuclear antibody test result. *Am J Med*. 2013 Apr;126(4):342-8. doi: 10.1016/j.amjmed.2012.09.014. Epub 2013 Feb 8. PMID: 23395534.
- Grygiel-Górniak B, Rogacka N, Puszczewicz M. Antinuclear antibodies in healthy people and non-rheumatic diseases - diagnostic and clinical implications. *Reumatologia*. 2018;56(4):243-248. doi:10.5114/reum.2018.77976
- Solomon DH, Kavanaugh AJ, Schur PH; American College of Rheumatology Ad Hoc Committee on Immunologic Testing Guidelines. Evidence-based guidelines for the use of immunologic tests: antinuclear antibody testing. *Arthritis Rheum*. 2002 Aug;47(4):434-44. doi: 10.1002/art.10561. PMID: 12209492.
- Satoh M, Chan EK, Ho LA, et al. Prevalence and sociodemographic correlates of antinuclear antibodies in the United States. *Arthritis Rheum*. 2012;64(7):2319-2327. doi:10.1002/art.34380
- Wandstrat AE, Carr-Johnson F, Branch V, Gray H, Fairhurst AM, Reimold A, Karp D, Wakeland EK, Olsen NJ. Autoantibody profiling to identify individuals at risk for systemic lupus erythematosus. *J Autoimmun*. 2006 Nov;27(3):153-60. doi: 10.1016/j.jaut.2006.09.001. Epub 2006 Oct 17. PMID: 17052888.

## Neuropsychiatric references

- de Amorim LC, Maia FM, Rodrigues CE. Stroke in systemic lupus erythematosus and antiphospholipid syndrome: risk factors, clinical manifestations, neuroimaging, and treatment. *Lupus*. 2017 Apr;26(5):529-536. doi: 10.1177/0961203316688784. PMID: 28394226.
- Jafri K, Patterson SL, Lanata C. Central Nervous System Manifestations of Systemic Lupus Erythematosus. *Rheum Dis Clin North Am*. 2017 Nov;43(4):531-545. doi: 10.1016/j.rdc.2017.06.003. Epub 2017 Aug 23. PMID: 29061240.
- Ricarte IF, Dutra LA, Abrantes FF, Toso FF, Barsottini OGP, Silva GS, de Souza AWS, Andrade D. Neurologic manifestations of antiphospholipid syndrome. *Lupus*. 2018 Aug;27(9):1404-1414. doi: 10.1177/0961203318776110. Epub 2018 May 17. PMID: 29768970.