



Obesity: Evaluation, treatment, & how to talk about it with patients

Ashleigh Pona, PhD
Assistant Professor - Clinical
Department of Psychiatry & Behavioral Health
The Ohio State University Wexner Medical Center

MedNet21
Center for Continuing Medical Education

 **THE OHIO STATE UNIVERSITY**
WEXNER MEDICAL CENTER

Agenda

- Background
- Evaluation
- Treatment
- Talking about weight with your patients

Background

Prevalence (2020)

- Body mass index of ≥ 30
- US adult obesity rate: 42.4%
 - First time national rate exceeded 40%
 - Lowest rate of obesity: Colorado, 23.6%
- US child (ages 2-19) obesity rate: 19.3%
 - Increasing with time, exhibiting earlier onset

Trust for America's Health, 2020

Causes

- Complex health issue with interacting, multifactorial causes
 - Obesogenic environment
 - Hereditary
 - Socioeconomic and sociocultural
 - Individual behaviors (physical activity, diet, medication use)

Hruby & Hu, 2015; Trust for America's Health, 2020

Demographic trends

- Protective factors
 - Higher income (vs lower)
 - More education (vs less)
 - Living in suburban & metro areas (vs rural)
- Risk factors
 - Poverty
 - Discrimination
 - Black 49.6%, Hispanic 44.8%, white 42.2%, Asian 17.4%
 - Food insecurity
 - Increased during COVID-19 crisis

Trust for America's Health, 2020

Consequences

- Associated with:
 - Poorer mental health, reduced quality of life
 - Leading causes of death: T2DM, heart disease, stroke, some types of cancer
 - More serious consequences of COVID infection, including hospitalization and death

Hruby & Hu, 2015; Trust for America's Health, 2020

Recommendations to address

- National organizations recommend physicians screen for obesity & provide intensive behavioral counseling
- However, obesity is not well managed in current health systems
 - Lack of training of healthcare workforce
 - Baseless assumptions of people with obesity
 - Lack of experience working in multidisciplinary teams
 - Lack of training in behavior change strategies

Campbell-Scherer et al., 2020

Role of primary care

- Small portion of adults with obesity ask a healthcare professional about weight loss
 - Of these, most consult their PCP
- Primary care: main point of contact for most people seeking health services
- Numerous articles have detailed strategies to manage obesity in primary care

Campbell-Scherer et al., 2020; Forgione et al., 2018

Evaluation

Evaluation

- 5As model for weight management counseling in primary care:
 - Assess
 - Advise
 - Agree
 - Assist
 - Arrange

Fitzpatrick et al., 2016; Vallis et al., 2013

Assess

- Screening for obesity, comorbidities, patient's willingness to make health behavior changes
- Using appropriate language without indication of stigma & shame
 - Patients prefer providers refer to their weight or BMI
 - Caution against the "personal responsibility" notion

Fitzpatrick et al., 2016

Assess

- BMI & waist circumference (visceral adiposity)
- Obesity-related complications
- History: diet, exercise, sleep, mental health, medications
- Characteristics & comorbidities associated with poor weight loss
 - Binge eating, sleep disorders, depression, chronic pain
 - Weight loss outcomes differ by race/ethnicity

Fitzpatrick et al., 2016; Forgione et al., 2018

Assess

- Readiness to change
 - Barriers: more pressing health or mental health issues, lacking self-efficacy, financial or psychosocial problems
 - If not ready: plan to address barriers, invite patient to inform you when ready, build on patient's confidence
 - If ready: praise efforts, what methods have been successful, ask how you can help, acknowledge their value of health

Fitzpatrick et al., 2016

Advise

- Counseling patient about:
 - Health risks of current weight
 - May influence patient's motivation
 - Health benefits of modest weight loss
 - Individualized diet plans & gradual change → long-term adherence

Fitzpatrick et al., 2016

Agree (on goals)

- Goal setting: key health behavior change strategy
 - SMART: Specific, Measurable, Attainable, Relevant, Time-based
 - Unrealistic goals can lead to failure & disappointment
- Collaborative approach
 - Initial weight loss goal of 5-10% of weight
- Self-monitoring, mobile applications

Fitzpatrick et al., 2016

Assist

- Problem solving: identifying barriers in achieving goals & developing plan with clear strategies to overcome
- ADAPT: Attitude (normalizing), Define problem, Alternative solutions, Predict consequences, Try out solution
- Some patients may require more intensive counseling
 - Consider referrals: behavioral psychologist, dietician, commercial programs

Fitzpatrick et al., 2016

Arrange

- Increase accountability through regular (e.g., monthly) follow-up
 - Assess patient's progress towards goals
 - Review self-monitoring records
 - Problem-solve barriers

Fitzpatrick et al., 2016

Treatment

Treatment

- Important to treat obesity as a **chronic, relapsing, multifactorial disease**
 - Nutrition, physical activity, emotion/behavior, medication
- Primary care counseling alone has limited ability to achieve clinically meaningful weight loss
- More benefit is seen with:
 - Added pharmacotherapy
 - Intensive counseling from dietitian or nurse + meal replacement therapy

Bronner, 2016; Tsai & Wadden, 2009

Nutrition

- Language matters
- Nutrition planning should be individualized
 - No one diet that is better than the rest
 - Depends on patient's motivation, resources, finances, personal preference
- Recommend 25% less calories
 - Not by restriction, but by improving calorie choices
- Focus on mindfulness of eating, including self-monitoring
- Viewing nutrition as a lifestyle change

Bronner, 2016; Taylor, 2020

Physical activity

- Language matters
- Physical activity planning should be individualized
 - Remember: SMART goals
- Get your patients moving – any movement helps
 - Get creative with ideas
- Self-monitoring can be helpful

Bronner, 2016

Emotion/Behavior

- Anxiety & depression are prevalent
 - Can impact eating behavior & adherence (decreased motivation)
- Screen for eating disorders
- Discuss alcohol & substance use
- Assess sleep
- Discuss eating habits
 - Dining out, distracted eating, stress eating, meal planning

Bronner, 2016

Medication

- Anti-obesity medications (& surgery) change the physiology of body regulation & offer best chance for long-term weight loss
 - Cannot replace diet, exercise, & lifestyle modification
 - May help to feel less hungry or full sooner, or make it harder for the body to absorb fat from foods
 - Numerous US FDA-approved medications currently available

Bronner, 2016

Medication

1. Identify medications possibly contributing to weight gain & change patient's regimen
2. Identify if patient meets FDA-approved anti-obesity medication indications
3. Trial medication
 - If no improvement after 3-4 months, consider different medication or increase dose
4. Medication as adjunct treatment

Bronner, 2016

Medication

Medication	How it works	Weight loss at 1 year
Orlistat	Works in gut to reduce amount of fat the body absorbs from food consumed	-5.5 lb with 60 mg -7.5 lb with 120 mg
Liraglutide	Mimics a hormone (glucagon-like peptide-1) that targets areas of brain that regulate appetite & food intake	-13.5 lb
Qsymia (phentermine & topiramate)	May lead to feeling less hungry or feeling full sooner	-14.5 lb with 7.5/46 mg -19.5 lb with 15/92 mg
Contrave (bupropion & naltrexone)	May lead to feeling less hungry or feeling full sooner	-13.5 lb

* Be aware of contraindications & adverse effects of different medications

Taylor, 2020

Bariatric surgery (referral)

- Promotes weight loss by restricting amount of food the stomach can hold, causing malabsorption of nutrients, or by a combination of both restriction & malabsorption
 - Does not replace diet, exercise, & lifestyle modification!

**Talking about weight
with your patients**

Weight bias & stigma

- Both healthcare professionals & patients with obesity endorse weight bias attitudes & beliefs about obesity
- Patients with obesity perceive biased treatment in healthcare, & this impacts how they access healthcare services
- Avoid making assumptions or judgments about patients' health & behaviors based on their weight

Campbell-Scherer et al., 2020

Weight bias & stigma

- Weight bias can be:
 - Subtle & overt
 - Verbal, physical, relational, cyber
- Can lead to rejection, prejudice, & discrimination
- Individuals affected may be:
 - Reluctant to seek medical care
 - Likely to delay important preventative healthcare services
 - Cancel medical appointments

Bronner, 2016

Weight bias & stigma

- Be aware of the following **misperceptions** of individuals with obesity:
 - Non-adherent
 - Dishonest
 - Lazy
 - Lacking in self-control
 - Unintelligent

Bronner, 2016; Puhl & Brownell, 2006

Obesity terminology

- **People-first language:** recognizes the potential dangers of labeling individuals by their disease
 - Say “patient with obesity” instead of “obese patient”
- Use **preferred/encouraged terms** & avoid discouraged terms

Encouraged	Discouraged
Weight Unhealthy weight Overweight Body mass index Affected by obesity	Morbidly obese Obese Fat

Bronner, 2016; Puhl et al., 2013

Considerations for clinic environment

- Ensure clinic furniture & equipment is appropriately sized for individuals with obesity
 - Chairs, toilets, doorways
 - Scales, gowns, blood pressure cuffs
- Ensure scales are in private areas
- Ensure staff are educated about obesity & weight bias

Kahan, 2018

Other weight bias reduction strategies

- Assess your own weight bias attitudes & beliefs
- Be mindful of patient's previous weight bias experiences & internalized weight bias
- Recognize & acknowledge multiple determinants of weight
- Separate weight from health – explore all causes of presenting problems
- Highlight importance of behavioral goals vs weight loss goals

Campbell-Scherer et al., 2020; Thille, 2019

References

- Bronner, T. G. (2016). *Introduction to Obesity Medicine: Helping Patients LOSE Weight and Helping Providers GAIN Professional Satisfaction* [PowerPoint slides]. Obesity Treatment Foundation. <https://www.hindsonfoundation.com/wconf/Bronner.pdf>
- Campbell-Scherer, D., Walji, S., Kemp, A., Piccinini-Vallis, H., & Vallis, T. M. (2020). Canadian Adult Obesity Clinical Practice Guidelines: Primary Care and Primary Healthcare in Obesity Management. Downloaded from: <https://obesitycanada.ca/guidelines/primarycare>
- Fitzpatrick, S. L., Wischenka, D., Appelhans, B. M., Pbert, L., Wang, M., Wilson, D. K., & Pagoto, S. L. (2016). An evidence-based guide for obesity treatment in primary care. *The American journal of medicine*, 129(1), 115-e1.
- Forgione, N., Deed, G., Kilov, G., & Rigas, G. (2018). Managing obesity in primary care: breaking down the barriers. *Advances in therapy*, 35(2), 191-198.
- Hruby, A., & Hu, F. B. (2015). The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*, 33(7), 673–689. <https://doi.org/10.1007/s40273-014-0243-x>
- Kahan, S. I. (2018, March). Practical strategies for engaging individuals with obesity in primary care. In *Mayo Clinic Proceedings* (Vol. 93, No. 3, pp. 351-359). Elsevier.
- Puhl, R., & Brownell, K.D. (2006). Confronting and coping with weight stigma: An investigation of overweight and obese individuals. *Obesity*, 14, 1802-1815.
- Puhl, R., Peterson, J. L., & Luedicke, J. (2013). Motivating or stigmatizing? Public perceptions of weight-related language used by health providers. *International journal of obesity*, 37(4), 612-619.
- Taylor, J. (2020). Looking beyond lifestyle: A comprehensive approach to the treatment of obesity in the primary care setting. *The Journal for Nurse Practitioners*, 16(1), 74-78.
- Thille, P. (2019). Managing anti-fat stigma in primary care: an observational study. *Health communication*, 34(8), 892-903.
- Trust for America's Health (2020). The state of obesity: Better policies for a healthier America. fah.org/stateofobesity2020
- Tsai, A. G., & Wadden, T. A. (2009). Treatment of obesity in primary care practice in the United States: A systematic review. *Journal of general internal medicine*, 24(9), 1073-1079.
- Vallis, M., Piccinini-Vallis, H., Sharma, A. M., & Freedhoff, Y. (2013). Modified 5 As: Minimal intervention for obesity counseling in primary care. *Canadian Family Physician*, 59(1), 27-31.