



## The Rise in Syphilis and the Role of the Emergency Department

**Ashley Lipps, MD**

*Assistant Professor, Division of Infectious Diseases  
Department of Internal Medicine  
The Ohio State University Wexner Medical Center*

MedNet21  
Center for Continuing Medical Education

THE OHIO STATE UNIVERSITY  
WEXNER MEDICAL CENTER

## Disclosures/Conflicts of Interest

- None

## Case Presentation

- 75 year old woman who presents to ED with progressive left vision loss.
- Symptoms started about 1 month prior; was seen at an outside facility where she was found to have left sided choroidal infarcts on exam – at the time she also endorsed headaches and was found to have elevated inflammatory markers – so the diagnosis of giant cell arteritis (GCA) was made
- Patient was treated with 1 gram IV methylprednisolone, followed by 75 mg daily of prednisone x 1 month (to present)

## Case Presentation

- At some point did undergo a temporal artery biopsy, which was negative for findings suggestive of GCA
- Reported that initially visual symptoms improved with steroids, but over the last several days she noticed significant decrease in vision in the left eye
- Ophthalmologic exam was notable for active choroiditis with new uveitis/vitritis in the left eye. Admitted to the hospital for further evaluation

### Case Presentation

- Further history reveals that the patient currently lives at home by herself in Ohio. She has 2 cats at home and no other animal exposures
- Currently retired (worked in retail in the past)
- No recent travel, no history of any international travel
- Has 2 adult children who live out of state
- No tobacco, alcohol or other drug use
- Not currently sexually active

### Case Presentation

- Physical exam was unremarkable other evidence of a very faint, healing rash on the trunk and upper arms
- On further questioning, the patient reports that several weeks ago she developed a severe rash over her entire body – went to an urgent care and was diagnosed with a bad allergic reaction. States the rash has been improving slowly over time.

### Case Presentation

- Patient reported she had a male new sexual partner about 6 months prior, although they are no longer in contact
- Barrier protection used but not every time
- She reports that the prior partner had several other sexual partners (both men and women); she was screened for HIV a few months ago after her partner notified her that he may have had unprotected sexual contact with a person with HIV
- No prior history of gonorrhea, chlamydia, syphilis or HSV

### Case Presentation

- HIV 1/2 Ab/p24 Ag: Non-reactive
- Urine/oral chlamydia/gonorrhea NAAT: Negative
- **Syphilis IgM/IgG: REACTIVE**
- **RPR: 1:512**
- Lumbar puncture: WBC 15, RBC <3, Protein 62, glucose 75
- CSF VDRL: Reactive 1:2

## Taking a sexual history

BOX 1. The Five P's approach for health care providers obtaining sexual histories: partners, practices, protection from sexually transmitted infections, past history of sexually transmitted infections, and pregnancy intention

1. **Partners**
    - "Are you currently having sex of any kind?"
    - "What is the gender(s) of your partner(s)?"
  2. **Practices**
    - "To understand any risks for sexually transmitted infections (STIs), I need to ask more specific questions about the kind of sex you have had recently:"
    - "What kind of sexual contact do you have or have you had?"
      - "Do you have vaginal sex, meaning 'penis in vagina' sex?"
      - "Do you have anal sex, meaning 'penis in rectum/anus' sex?"
      - "Do you have oral sex, meaning 'mouth on penis/vagina'?"
  3. **Protection from STIs**
    - "Do you and your partner(s) discuss prevention of STIs and human immunodeficiency virus (HIV)?"
    - "Do you and your partner(s) discuss getting tested?"
    - For condoms:
      - "What protection methods do you use? In what situations do you use condoms?"
  4. **Past history of STIs**
    - "Have you ever been tested for STIs and HIV?"
    - "Have you ever been diagnosed with an STI in the past?"
    - "Have any of your partners had an STI?"
- Additional questions for identifying HIV and viral hepatitis risks**
- "Have you or any of your partner(s) ever injected drugs?"
  - "Is there anything about your sexual health that you have questions about?"
5. **Pregnancy intention**
    - "Do you think you would like to have (more) children in the future?"
    - "How important is it to you to prevent pregnancy (until then)?"
    - "Are you or your partner using contraception or practicing any form of birth control?"
    - "Would you like to talk about ways to prevent pregnancy?"

Source: CDC Sexually Transmitted Infections Treatment Guidelines, 2021

## Taking a sexual history

- Establish rapport and make your patient feel comfortable before asking sensitive questions
- Use neutral and inclusive terms (e.g. partner) and pose your questions in a non-judgmental manner
- Avoid making assumptions about your patients' sexual orientation, gender identity or sexual behaviors based on age, appearance, marital status, or other factors


## Syphilis: a review



- Syphilis is caused by the spirochete *Treponema pallidum*
- Major mode of transmission is via sexual contact
- Vertical transmission can occur (congenital syphilis)
- Can cause a wide variety of clinical manifestations, including periods of clinical latency (asymptomatic) if left untreated

## Syphilis: a review

Primary and Secondary Syphilis — Rates of Reported Cases Sex, United States, 2010–2019



Source: U.S. Centers for Disease Control and Prevention

## Syphilis: a review


Primary and Secondary Syphilis — Distribution of Cases by Sex and Sex of Sex Partners, United States, 2019



Source: U.S. Centers for Disease Control and Prevention

## Congenital Syphilis


Congenital Syphilis — Reported Cases by Vital Status and Clinical Signs and Symptoms\* of Infection, United States, 2015–2019



Source: U.S. Centers for Disease Control and Prevention

## Congenital Syphilis

Congenital Syphilis — Missed Prevention Opportunities Among Mothers Delivering Infants with Congenital Syphilis, United States, 2015–2019




Source: U.S. Centers for Disease Control and Prevention

## Primary syphilis

- Painless ulcer (chancre) appears at site of inoculation – can go unnoticed depending on the location.
- Regional lymphadenopathy can occur (inguinal, cervical)
- Chancres are highly infectious and may resolve without treatment within 1-6 weeks

## Primary syphilis



Source: Centers for Disease Control and Prevention Public Health Image Library

## Secondary syphilis


- Typically occurs about 4-8 weeks after onset of primary chancre, more likely to prompt medical evaluation
- The classic symptom is a diffuse maculopapular rash, which commonly involves the palms, soles, chest and back
- Lymphadenopathy, malaise, fever, mucous patches (genitals, mouth), patchy alopecia, and condyloma lata can occur as well

## Secondary syphilis



Source: Negusse Ocbamichael, PA; Public Health—Seattle & King County STD Clinic

## Secondary syphilis




Source: Negusse Ocbamichael, PA; Public Health—Seattle & King County STD Clinic  
Source: Centers for Disease Control and Prevention Public Health Image Library

## Tertiary Syphilis

- Form of late syphilis – can occur decades after initial infection if treatment is not administered
- Gummatous disease (granulomatous disease of skin, subcutaneous tissues, bones or viscera)
- Cardiovascular syphilis (involvement of vasa vasorum – aortic aneurysm, aortic insufficiency)

## Tertiary Syphilis



Source: Centers for Disease Control and Prevention Public Health Image Library

## Latent syphilis

- Early latent syphilis (infection of less than 1 year duration)
- Late latent syphilis (infection greater than 1 year duration)
- Latent syphilis of unknown duration

## Neurosyphilis, ocular syphilis, otosyphilis

- CNS involvement can occur during any stage of infection
- Early neurosyphilis - cranial nerve dysfunction, meningitis, meningovascular syphilis, stroke and/or acute altered mental status
- Late neurosyphilis – general paresis/tabs dorsalis (less common)
- Ocular syphilis (anterior, posterior or pan-uveitis), can occur with or without other associated neurologic manifestations
- Otosyphilis: usually presents with tinnitus, vertigo, sensorineural hearing loss

## Screening for syphilis


Syphilis	
Women and Men	<ul style="list-style-type: none"> <li>Screen asymptomatic adults at increased risk (history of incarceration or commercial sex work, geography, race/ethnicity, and being a male younger than 29 year) for syphilis infection<sup>2,7</sup></li> </ul>
Pregnant Women	<ul style="list-style-type: none"> <li>All pregnant women at the first prenatal visit<sup>8</sup></li> <li>Retest at 28 weeks gestation and at delivery if at high risk (lives in a community with high syphilis morbidity or is at risk for syphilis acquisition during pregnancy (using drugs, STIs during pregnancy, multiple partners, a new partner, partner with STIs)<sup>2</sup></li> </ul>
Men Who Have Sex With Men (MSM)	<ul style="list-style-type: none"> <li>At least annually for sexually active MSM<sup>2</sup></li> <li>Every 3 to 6 months if at increased risk<sup>2</sup></li> </ul>
Transgender and Gender Diverse People	<ul style="list-style-type: none"> <li>Consider screening at least annually based on reported sexual behaviors and exposure<sup>2</sup></li> </ul>
Persons with HIV	<ul style="list-style-type: none"> <li>For sexually active individuals, screen at first HIV evaluation, and at least annually thereafter<sup>9,10</sup></li> <li>More frequent screening might be appropriate depending on individual risk behaviors and the local epidemiology<sup>2</sup></li> </ul>

Source: CDC Sexually Transmitted Infections Treatment Guidelines, 2021

## Screening for syphilis

- Laboratory testing - Reverse sequence algorithm
- Treponemal specific tests: Syphilis IgM/IgG, T.pallidum particle agglutination assay (TP-PA)
- Non-treponemal specific tests: rapid plasma reagin (RPR)

## Diagnosis



## Treatment

- Early syphilis (primary, secondary, early latent): 2.4 million units Benzathine penicillin G IM in a single dose
  - Alternative for penicillin allergic, non-pregnant adults: doxycycline 100 mg twice daily x 14 days

## Treatment

- Late syphilis (late latent syphilis, latent syphilis of unknown duration, tertiary syphilis if CNS disease excluded): 2.4 million units Benzathine penicillin G IM weekly x 3 doses
  - Alternative for penicillin allergic, non-pregnant adults: doxycycline 100 mg twice daily x 28 days

## Treatment

- Neurosyphilis: Aqueous crystalline penicillin G 18-24 million units per day, administered as 3-4 MU IV every 4 hours or continuous infusion given for 10-14 days
  - Alternative procaine penicillin G 2.4 million units IM once daily plus probenecid 500 mg orally 4 times/day for 10-14 days

## Other treatment considerations

- All people with syphilis should be screened for HIV
- Syphilis exposure has been associated with an increased risk of future HIV acquisition, particularly in men—counseling on safer sex practices and HIV Pre-Exposure Prophylaxis (PrEP)
- Sexual partners should be treated

Peterman TA, Newman DR, Maddox L, Schmitt K, Shiver S. High risk for HIV following syphilis diagnosis among men in Florida, 2000-2011. *Public Health Rep.* 2014;129(2):164-169.

## References

- Workowski K, Bachmann L, Chan P, et al. *Sexually Transmitted Infections Treatment Guidelines*, 2021. MMWR Recomm Rep 2021; 70. <https://www.cdc.gov/std/treatment-guidelines/default.htm>. Accessed 8-12-21
- Centers for Diseases Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention – Syphilis Statistics, 2021. <https://www.cdc.gov/std/syphilis/stats.htm>. Accessed 8-12-21
- National Coalition for Sexual Health. Sexual Health and Your Patients: A Provider's Guide. <https://nationalcoalitionforsexualhealth.org/tools/for-healthcare-providers/sexual-health-and-your-patients-a-providers-guide>. Accessed 8-12-21
- Peterman TA, Newman DR, Maddox L, Schmitt K, Shiver S. High risk for HIV following syphilis diagnosis among men in Florida, 2000-2011. *Public Health Rep.* 2014;129(2):164-169.



## Syphilis and co-existent sexually transmitted diseases

## Syphilis re-infection?



### Sexually Transmitted Diseases

**Sommer E. Lindsey, MD, FACEP**

*Assistant Professor*

*Department of Emergency Medicine*

*The Ohio State University Wexner Medical Center*

**MedNet21**  
Continuing Medical Education

**THE OHIO STATE UNIVERSITY**  
WEXNER MEDICAL CENTER

## Background

### National Data


- Sexually transmitted infections are on the rise
- **Syphilis**
  - 35,063 new cases since 2014, 71% increase
  - Several demographics have shown increased case numbers
  - Gonorrhea and syphilis increase the likelihood of transmission of HIV

## STI testing in Urban Emergency Departments

### Ideal Population

- Indigent population
- Uninsured
- No Primary Care
- ED is point of healthcare access
- High risk populations:
  - Minorities
  - Transient/homeless
  - IVDU
  - prostitution
  - multiple partners with diverse sexual orientation
- Perfect opportunity to screen for syphilis in a population that is under tested and under treated

## Primary and Secondary Syphilis — Rates of Reported Cases by State and Territory, United States, 2018



## Franklin County

- Ranks 21<sup>st</sup> amongst counties in nation in number of new cases of syphilis
- Half of all new cases of syphilis in just 28 counties nationally
  - Less than 1% counties nationwide
- One of 48 counties identified nationally as HIV hot spot

Source: CDC


## Emergency department visits: trends

YEAR	NHAMCS ESTIMATED ED VISITS (MILLIONS)
2001	107.5
2002	110.2
2003	113.9
2004	110.2
2005	115.3
2006	119.2
2007	116.8
2008	123.8
2009	136.1
2010	129.8
2011	136.3
2012	130.9
2013	130.4
2014	141.4
2015	135.9
2016	145.6

Source: CDC

Patena W, Peterson T, Gish T. An Increase in Sexually Transmitted Infections from US Emergency Departments. Prev Med. 2017 July; 98:143-146. <https://doi.org/10.1016/j.ypmed.2017.05.007>

### Order practice prior to initiation of study



### Why didn't we do this sooner?

- It's Complicated



Source: CDC

<https://www.jfpe.org/why-didnt-we-test-earlier-for-syphilis-in-the-united-states/>  
JAMA Network Open. 2019;2(6):e19042. doi:10.1001/jamanetworkopen.2019.5042 (Reprinted) June 11, 2019 2/4

### Why didn't we do this sooner?

- Practitioners unaware of syphilis surge
- Who is responsible for follow-up on these results?
- Interpretation of results
- Tracking patients down
- Collaboration with outpatient clinics
- Linkage to care and initiation of PrEP
- Insurance coverage: US Preventative Services Task Force
  - Medicaid mostly cover routine screening or “medically necessary” testing



Source: CDC

<https://www.jfpe.org/why-didnt-we-test-earlier-for-syphilis-in-the-united-states/>  
JAMA Network Open. 2019;2(6):e19042. doi:10.1001/jamanetworkopen.2019.5042 (Reprinted) June 11, 2019 2/4

### The test has been there. Why aren't you ordering it?

- Survey of all EM faculty, residents, NPs at OSU Main and East
- Questions address hesitation to ordering HIV/syphilis testing from ED

Source: CDC



### Solution: Guarantee follow-up outside the ED

- ID generated list daily
  - All patients tested for HIV/syphilis and their results
- Interpretation of results
- Contacting patient
  - CPH and ODH help
- Arranging for treatment or continued surveillance
- PrEP
- STI ID Attending on call pager on WebExchange

Carter MW, Wu H, Cohen S, et al. Linkage and referral to HIV and other medical and social services: a focused literature review for sexually transmitted disease prevention and control programs. Sex Transm Dis. 2016;43(2 Suppl 1):S76-S82. [Systematic review, 33 studies]

### ED STI Protocol      Management of STIs

**Patient examination:**  
Previous STIs including HIV status, barrier protection, sex of partners, number of partners, rectal, oral, vaginal intercourse

#### STI orders:

**Female:** gonorrhea and chlamydia (cervix)  
Affirm (wet prep-BV, yeast, trich)  
Syphilis (STAT not next day lab)  
Rapid HIV (blood)

**Male:** gonorrhea and chlamydia (urine/urethra)  
Urine micro (trich)  
Syphilis (STAT not next day lab)  
Rapid HIV (blood)

## ED STI Protocol      Management of STIs

May provide presumptive treatment for gonorrhea and chlamydia based upon history, exam, and/or high-risk status while in the ED.

If HIV or syphilis +, notification will be sent to Infectious Disease and the health department for follow up (by lab, not the ED provider). ID will then contact the patient regarding treatment options/locations, follow-up, and PrEP initiation for high risk patients.

## ED STI Protocol      Management of STIs

**Current CDC treatment Recommendations:**  
**Chlamydia:** Zithromax 1000 mg PO x1 OR Doxycycline 100mg BID x7 days  
**Gonorrhea:** Rocephin 250mg IM x1 PLUS Zithromax 1000 mg PO x1  
**Trichomoniasis:** Flagyl 2000 mg PO x1 OR Flagyl 500mg BID x7 days  
**Syphilis:** Benzathine (PCN G) 2.4 million units IM x1

Update: Gonorrhea 500mg IM x 1 for 300 lbs or less, 1 gram IM x 1 for greater than 300 lbs

## ED protocol

- STI-related complaint/Concern for STI based on clinical presentation
- History
  - Number sexual partners
  - Known HIV or syphilis diagnosis?
  - Barrier methods used
  - Sexual contact
    - Need for oral, rectal, and/or vaginal swabs
- Test for GC/Chlamydia (oral, urine, rectal, urethral, vaginal swabs), HIV (serum), syphilis (serum)
- Rapid HIV, with p24 antigen and syphilis AB with reflex RPR

Source: CDC

## ED Order Set

UPDATE: Rapid Gonorrhea and Chlamydia testing now available. ALSO added pregnancy test and UA to streamline ordering

- Order set
  - Type “STI” in order set box on IHIS


## STI Labs &amp; Medications Panel

Please select gender appropriate orders for STI screening	
<input checked="" type="checkbox"/> CHLAM & GONORRHEA: AMP CERVIX	STAT, ONE TIME For 1 Occurrences CERVIX
<input checked="" type="checkbox"/> VAGINITIS DNA PROBES	URGENT, ONE TIME For 1 Occurrences VAGINA
<input type="checkbox"/> CHLAM & GONORRHEA: AMP, URINE	STAT, ONE TIME For 1 Occurrences URINE FIRST CATCH
<input type="checkbox"/> CHLAM & GONORRHEA: AMP, ORAL	STAT, ONE TIME For 1 Occurrences THROAT
<input type="checkbox"/> CHLAM & GONORRHEA: AMP, RECTAL	STAT, ONE TIME For 1 Occurrences RECTUM
<input checked="" type="checkbox"/> SYPHILIS AB W/REFLEX RPR	STAT, ONE TIME For 1 Occurrences
<input checked="" type="checkbox"/> RAPID HIV-1/HIV-2 AB WITH P24 ANTIGEN	STAT, ONE TIME For 1 Occurrences
<input type="checkbox"/> HIV VIRAL LOAD RNA PCR QUANT (Use in addition to rapid HIV test if acute HIV suspected)	STAT, ONE TIME For 1 Occurrences
<input checked="" type="checkbox"/> azithromycin (ZITHROMAX) tablet	1,000 mg, Oral, ONCE For 1 Doses
<input checked="" type="checkbox"/> ceftriaxone (ROCEPHIN) injection	250 mg, Intramuscular, ONCE For 1 Doses
<input type="checkbox"/> metronidazole (FLAGYL) tablet	2,000 mg, Oral, ONCE For 1 Doses
<input type="checkbox"/> ondansetron (ZOFTRAN-ODT) disintegrating tablet	4 mg, Oral, ONCE For 1 Doses

Dosing has been changed


Source: CDC

## Data since initiation of project




Source: CDC

## STI Testing in the OSU East Emergency Department 2012-2017



54

## HIV and syphilis tests ordered from OSU EDs



## SYPHILIS

- Between Nov 1 2018 and Nov 30 2019 there were **57 positive syphilis antibody tests**
  - 24 positive tests in women
  - 33 positive tests in men
- Totals:
  - 27/57 Previously treated infections
  - 19/57 Late latent infections (6 fully treated, 4 partially treated, 9 untreated)
  - 2/57 Secondary syphilis (2/2 fully treated)
  - 1/57 Primary syphilis (1/1 fully treated)
  - 8/57 false positives
  - 16 positive and/or inadequately treated cases found
    - 1 % of those tested had a positive result and inadequate/no treatment

Source: CDC

## Moving Forward

- Protocol for STI testing in EDs nationally
- Exemplar of interdepartmental collaboration with OSU Infectious Disease and collaboration with Columbus Public Health
- Model for quick linkage to care and initiation of PrEP
  - PrEP can reduce risk of HIV acquisition through sex by 90%
  - Navigators in ED who will assist patients with LTC and PrEP
- Social Work resources
- Nurse case manager, establish primary care

Source: CDC

## Goals

- PrEP referral in STI order set to specific sites
  - ID clinic, THW, FACES, Equitas, primary care, patient choice
- HPV vaccine in appropriate patients
- Introduce model to other area healthcare systems
- Retrospective analysis of how early detection/treatment of HIV reduces number of ED visits/year
- Study demographics of patient populations being tested
  - Visits to ED/year
  - Race
  - Age
  - Gender
  - Sexual Orientation
  - Insurance status

Source: CDC