


# **Adrenal Insufficiency: Current Practice 2012**

**Lawrence S. Kirschner, MD, PhD**  
**Professor of Medicine**  
**Division of Endocrinology, Diabetes, and Metabolism**  
**The Ohio State University's Wexner Medical Center**


## **Overview**

- **A very brief review of adrenal function**
- **What is adrenal insufficiency?**
- **Adrenal insufficiency in the outpatient setting**
- **Adrenal insufficiency during critical illness**
- **Therapy for adrenal insufficiency**

# Hypothalamus-Pituitary-Adrenal Axis



# Actions of cortisol



- **Actions of Aldosterone**
  - Promotes sodium/water retention
  - Promotes potassium excretion
  - May be involved in tissue remodeling (e.g. in the heart)
- **Actions of adrenal androgens**
  - Responsible for initiation of puberty
    - Secondary sex characteristics in women

## **Adrenal Insufficiency (Addison disease)**




Image courtesy of  
Wellcome Images  
<http://images.wellcome.ac.uk/>

## What is Adrenal insufficiency?

- When discussing adrenal insufficiency (Addison disease), we are almost always talking about glucocorticoid (cortisol) insufficiency
- However, other adrenal hormones can also be affected in *primary* adrenal failure


## Clinical Addison disease



## Clinical Features of Chronic Adrenocortical Insufficiency


•Weakness, fatigue	100%
•Weight loss	100%
•Anorexia	100%
•Hyperpigmentation	92%
•Hypotension	88%
•Nausea, abdominal pain	56%
•Salt craving	19%
•Hypoglycemia	??
»more common in children and women	

Hyperpigmentation,  
toxic appearance



Images courtesy of  
Wellcome Images  
<http://images.wellcome.ac.uk/>

## Addison disease



Hyperpigmentation, including  
knuckles and palmar creases

### **Features of Acute Adrenocortical Insufficiency (Adrenal Crisis)**


- Hypotension
- Weakness (prox. muscle),  
confusion
- Nausea, vomiting, abdominal pain
- Dehydration, hypovolemia
- Hyperthermia
- Hypoglycemia

**TREAT FIRST, AND DIAGNOSE LATER!!**


### **Adrenal Crisis**

- **Acute loss of adrenal function**
  - Acute loss of adrenals
    - Surgery
    - Hemorrhage/thrombosis
  - Acute loss of pituitary function
  - Acute loss of steroid replacement
- OR
- **Acute stress in the setting of compensated chronic adrenal failure**
  - Precipitating event (e.g., like DKA)

## Normal Adrenal Function




## Primary Adrenal Insufficiency



- **Primary**
  - Adrenal gland
  - Destruction of glands

## Secondary Adrenal Insufficiency



- **Primary**
  - Adrenal gland
  - Destruction of glands
- **Secondary**
  - Pituitary
  - Inadequate ACTH
  - NO increased pigment
  - Fewer electrolyte imbalances

## Causes of adrenal failure

- Like CS, iatrogenic causes are probably most common
- **Inherited forms of adrenal failure**
  - Typically presenting early in life (<1 yr)
    - CAH, especially salt-wasters (steroid biosynthesis defect)
    - Other rare genetic diseases (lipoid CAH, AHC)
  - Typically presenting in childhood, and dx should be “obvious”
    - AAA
      - Alacrima, Achalasia, Adrenal failure
    - Autoimmune Polyendocrine Syndrome (APS), Type I (APECED)
      - Ectodermal dysplasia, mucocutaneous candidiasis



## Causes of adrenal failure

- Like CS, iatrogenic causes are probably most common
- Inherited forms of adrenal failure
  - Typically presenting early in life (<1 yr)
    - CAH, especially salt-wasters (steroid biosynthesis defect)
    - Other rare genetic diseases (lipoid CAH, AHC)
  - Typically presenting in childhood, and dx should be “obvious”
    - AAA
      - Alacrima, Achalasia, Adrenal failure
    - Autoimmune Polyendocrine Syndrome (APS), Type I (APECED)
      - Ectodermal dysplasia, mucocutaneous candidiasis


## Causes of adrenal failure

- Like CS, iatrogenic causes are probably most common
- Inherited forms of adrenal failure
  - Typically presenting early in life (<1 yr)
    - CAH, especially salt-wasters (steroid biosynthesis defect)
    - Other rare genetic diseases (lipoid CAH, AHC)
  - Typically presenting in childhood, and dx should be “obvious”
    - AAA
      - Alacrima, Achalasia, Adrenal failure
    - Autoimmune Polyendocrine Syndrome (APS), Type I (APECED)
      - Ectodermal dysplasia, mucocutaneous candidiasis

# Causes of adrenal failure


- Like CS, iatrogenic causes are probably most common
- Inherited forms of adrenal failure
  - Typically presenting early in life (<1 yr)
    - CAH, especially salt-wasters (steroid biosynthesis defect)
    - Other rare genetic diseases (lipoid CAH, AHC)
  - Typically presenting in childhood, and dx should be “obvious”
    - AAA
      - Alacrima, Achalasia, Adrenal failure
    - Autoimmune Polyendocrine Syndrome (APS), Type I (APECED)
      - Ectodermal dysplasia, mucocutaneous candidiasis

- Presenting later in life
  - Autoimmune Polyendocrine Syndrome (APS), Type II
    - Type I DM, thyroid disease
    - May occur as sole autoimmune feature (although rare)
  - Adrenal hemorrhage
    - Resulting from sepsis
  - HIV, other viral diseases
  - Adrenalectomy
- Note that non-classical CAH rarely causes adrenal insufficiency




## **Secondary adrenal failure**

- **Pituitary malfunction**
  - Tumor destroying normal cells
  - Autoimmune hypophysitis
    - May be quite specific for loss of ACTH-producing cells
  - Infiltrative diseases of pituitary
    - Histiocytosis X
    - Sarcoidosis
    - Metastatic disease



## Diagnosis of Adrenal Insufficiency in the Outpatient setting: Static Testing

- **A GOOD HISTORY IS ESSENTIAL!**
  - History of steroid use, including nasal steroids or injected steroids (e.g., back injections)
- **8 AM cortisol (probably NOT reliable in hospitalized patients)**




## Diagnosis of Adrenal Insufficiency in the Outpatient setting: Static Testing - 2

- ACTH measurements
  - Generally not helpful, particularly low values
  - Elevated values may suggest primary Adrenal Insufficiency in the right clinical setting
- “Suggestive” findings:
  - Eosinophilia, hyperchloremia, acidosis, hypercalcemia, azotemia, hyponatremia/hyperkalemia and fasting hypoglycemia

## Diagnosis of Adrenal Insufficiency: ACTH stim test

- Give IV/IM bolus of 250 mcg ACTH, measure blood at 0, 30, 60 min
- Normal response is for cortisol to reach >18 mcg/dl
- Caveat :ACTH stim test will be “normal” in early pituitary failure. Once adrenal atrophy sets in, test becomes subnormal




# **Adrenal Insufficiency: Current Practice 2012**


**Rami N. Khayat, MD**  
Associate Professor  
Pulmonary, Critical Care, and Sleep Medicine  
The Ohio State University's Wexner Medical Center

## **Adrenal Insufficiency during Critical Illness**

- **Adrenal function during critical illness**
- **Relative adrenal insufficiency**
- **Overview of Corticosteroid therapy in the ICU**
- **Conclusions**



## The Adrenal Response to Prolonged Critical Illness



*J Clin End Met.* 1995;80:1238.

## Adrenal Function in Critical Illness

- Adrenal Gland
  - Synthetic inhibition
    - Drugs
      - Etomidate
      - Ketoconazole
      - » Corticosteroids
    - Cytokines
  - Destruction
    - Pre-existing
      - Autoimmune
      - Infection
        - HIV • CMV
        - TB • Fungal
      - Metastasis
    - Acute
      - Hemorrhage
      - Infection

## Adrenal Function in Critical Illness

- ↑ Hepatic metabolism of cortisol
  - Rifampin
  - Phenytoin
  - Phenobarbital
- Glucocorticoid Resistance



# The Canadian Medical Association Journal

Vol. 43

TORONTO, JULY, 1940


No. 1

## ON THE THERAPEUTIC VALUE OF ADRENAL CORTICAL HORMONES IN TRAUMATIC SHOCK AND ALLIED CONDITIONS\*

BY HANS SELYE, CHRISTIANNE DOSNE, LUCY BASSETT AND JOAN WHITTAKER

**“The fact that cortical hormone therapy exerts beneficial effects in so many conditions makes it rather likely that the hormone is not a specific antidote in any one of these cases but raises shock resistance in general because a condition of relative adrenal insufficiency exists in organisms exposed to non-specific damage.”**

## Cortisol Levels-Marker of Survival




Venkatesh and Cohen 2011

# Cortisol and Septic Shock

%	Basal cortisol	$\Delta$ max	mortality
---	----------------	--------------	-----------

Annane JAMA 2000; 283:1038-1045

## Prognostic Value of Cortisol Levels and ACTH Response



Annane, D. et al. JAMA 2000;283:1038-1045

# JAMA

The Journal of the American Medical Association

Copyright 2002 by the American Medical Association. All Rights Reserved. Applicable FARS/DFARS Restrictions Apply to Government Use. American Medical Association, 515 N. State St, Chicago, IL 60610.

Volume 288(7)

21 August 2002

p 862-871

## Effect of Treatment With Low Doses of Hydrocortisone and Fludrocortisone on Mortality in Patients With Septic Shock

[Caring for the Critically Ill Patient]

Annane, Djillali MD, PhD; Sébille, Véronique PhD; Charpentier, Claire MD; Bollaert, Pierre-Edouard MD, PhD; François, Bruno MD; Korach, Jean-Michel MD; Capellier, Gilles MD, PhD; Cohen, Yves MD, PhD; Azoulay, Elie MD; Troché, Gilles MD; Chaumet-Riffaut, Philippe MD; Bellissant, Eric MD, PhD

## Steroids In Septic Shock

Catecholamine dependent septic shock (300)

ACTH Stim Test

76% Non-Responders (229)

24% Responders (70)

<9 ug/dl cortisol

>9 ug/dl cortisol

Steroids

Placebo

Steroids


Placebo

50 mg Hydrocortisone q6<sup>o</sup>  
50 ug Fludrocortisone qd

50 mg Hydrocortisone q6<sup>o</sup>  
50 ug Fludrocortisone qd


Annane JAMA 2002; 288:862-871

## Effect of Low Dose Hydrocortisone on Mortality in Patients with Septic Shock



Annane, D. et al. JAMA 2002;288:862-871

## Effect of Low Dose Hydrocortisone on Mortality in Patients with Septic Shock



Annane, D. et al. JAMA 2002;288:862-871

# Steroids In Septic Shock

## Mortality

28 Day

ICU

Hospital

1 Year

## Vasopressor Withdrawal

(28 days)

## Non- Responders

Steroids

Placebo

## Responders

Steroids

Placebo

53%\*

63%

61%

53%

58%\*

70%

67%

59%

61%\*

72%

69%

59%

68%\*

77%

69%

71%

57%\*

40%

50%

53%

Median

Median

Median

Median

7


10

9

7

Annane JAMA 2002; 288:862-871


## Effects of Corticosteroids on Mortality ICU Severe Sepsis and Septic Shock



bmj.com

Annane BMJ 2004;329:480

## Approach to Suspected Adrenal Insufficiency



## Concerns

- High mortality in the Control group
- Use of Etomidate
- Design and power
- Severe refractory shock required for enrollment

# CORTICUS

## *The* NEW ENGLAND JOURNAL *of* MEDICINE

ESTABLISHED IN 1812


JANUARY 10, 2008

VOL. 358 NO. 2

### Hydrocortisone Therapy for Patients with Septic Shock

Charles L. Sprung, M.D., Djillali Annane, M.D., Ph.D., Didier Keh, M.D., Rui Moreno, M.D., Ph.D., Mervyn Singer, M.D., F.R.C.P., Klaus Freivogel, Ph.D., Yoram G. Weiss, M.D., Julie Benbenishty, R.N., Armin Kalenka, M.D., Helmuth Forst, M.D., Ph.D., Pierre-Francois Laterre, M.D., Konrad Reinhart, M.D., Brian H. Cuthbertson, M.D., Didier Payen, M.D., Ph.D., and Josef Briegel, M.D., Ph.D., for the CORTICUS Study Group\*


### Hydrocortisone Therapy for Patients with Septic Shock Sprung et al NEJM 2008



#### Enrolment and Outcome

## Hydrocortisone Therapy for Patients with Septic Shock


### Kaplan–Meier curves for survival at 28 days according to response to corticotropin test



Sprung et al NEJM 2008

## Hydrocortisone Therapy for Patients with Septic Shock


### Kaplan–Meier Curves for Survival at 28 Days



Sprung et al NEJM 2008



# Corticosteroid Treatment and Intensive Insulin Therapy for Septic Shock in Adults




## Corticosteroid Treatment and Intensive Insulin Therapy for Septic Shock in Adults

Variables	Intensive Insulin Therapy (n = 255)	Conventional Glucose Control (n = 254)	P Value		Hydrocortisone + Fludrocortisone (n = 245)	Hydrocortisone Alone (n = 264)	P Value	
			Unadjusted	Adjusted <sup>a</sup>			Unadjusted	Adjusted <sup>a</sup>
In-hospital death, No./total (%)	117/255 (45.9)	109/254 (42.9)	.50	.37	105 (42.9)	121 (45.8)	.50	.91
Overall survival								
Deaths, No. (%)	122 (47.9)	118 (46.5)			112 (45.7)	128 (48.5)		
Kaplan-Meier estimate of survival rates, HR (95% CI), d	1.04 (0.80-1.34)	1 [Reference]	.78	.39	0.94 (0.73-1.21)	1 [Reference]	.61	.67
28	62.2 (56.4-68.5)	61.1 (55.3-67.5)			62.5 (56.6-68.9)	60.9 (55.2-67.1)		
90	51.8 (45.9-58.4)	54.8 (48.9-61.4)			54.2 (48.2-61.0)	52.4 (46.6-58.9)		
180	50.9 (45.0-57.6)	52.1 (46.2-58.8)			52.9 (46.9-59.7)	50.2 (44.4-56.8)		
No. of patients who died	103	82			105	121		
Causes of death, No. (%)								
Multiple organ failure	92 (78.6)	66 (60.6)	.004 <sup>b</sup>	.005 <sup>b</sup>	75 (71.4)	83 (68.6)	.67 <sup>b</sup>	.74 <sup>b</sup>
Cardiovascular	9 (8.7)	7 (6.5)			7 (6.7)	9 (7.4)		
Stroke	1 (1.0)	2 (2.4)			3 (2.9)	0		
Brain hemorrhage	0	2 (2.4)			0	2 (1.7)		
Refractory hypoxia	1 (1.0)	2 (2.4)			2 (1.9)	1 (0.8)		
Unknown	0	3 (3.7)			3 (2.9)	0		
No. of days, median (IQR)								
Vasopressor-free within the first 7 days	4 (1-0)	4 (2-5)	.58	.00	4 (2-5)	4 (1-5)	.02	.61
Mechanical ventilation-free within 28 days	10 (2-22)	13 (2-23)	.51	.29	12 (2-23)	12 (2-22.5)	.50	.81
Cumulative incidence of SOFA <8 at day 7 (95% CI)	64.3 (58.6-70.1)	60.6 (54.7-66.6)	.38	.75	63.3 (57.3-69.2)	61.7 (56.0-67.5)	.75	.78
Length of stay, median (IQR), d								
ICU								
All patients	9 (4-19)	9 (4-15)	.70	.39	9 (4-16)	9 (4-17.5)	.86	.35
Survivors	10 (6-19)	9 (5-15)	.68	.46	10 (6-16)	9 (5-17)	.52	.10
Hospital								
All patients	16 (6-34)	15 (7-30)	.87	.94	14 (6-25)	18 (7-34)	.15	.07
Survivors	24 (12-43)	22 (11-39)	.87	.57	19 (5-40)	25.5 (14-42)	.09	.13

**JAMA 2010;303:341-348**

## Corticosteroids for ARDS NEJM 2006 “ARDSNET”



**Figure 2.** Probability of Survival and the Proportion of Patients with Persistent ARDS Who Became Able to Breathe without Assistance during the First 180 Days after Randomization.

At 180 days, 29 patients in the placebo group had died, 58 had been discharged home, and 4 had not been discharged home; the respective values in the methylprednisolone group were 28, 57, and 4. The status was known for all 180 patients at 180 days.

## Role of Steroids in Specific Conditions

### Good

- Meningitis
- Typhoid fever
- Spinal cord injury
- Pneumocystis Carinii Pneumonia
- No evidence
  - Fibroproliferative ARDS?
  - Sepsis?

### Bad

Industrial strength  
pharmacologic  
does

## **Surviving Sepsis 2008**

- We suggest that intravenous hydrocortisone be given only to adult septic shock patients after it has been confirmed that their blood pressure is poorly responsive to fluid resuscitation and vasopressor therapy (grade 2C).
- We suggest that the ACTH stimulation test not be used to identify the subset of adults with septic shock who should receive hydrocortisone (grade 2B).
- We suggest that patients with septic shock should not receive dexamethasone if hydrocortisone is available (grade 2B).

## **Surviving Sepsis 2008**

- daily addition of oral fludrocortisone (50 µg) if hydrocortisone is not available and the steroid that is substituted has no significant mineralocorticoid activity. Fludrocortisone is considered optional if hydrocortisone is used (grade 2C).
- that clinicians wean the patient from steroid therapy when vasopressors are no longer required (grade 2D).
- We recommend that doses of corticosteroids comparable to >300 mg hydrocortisone daily not be used in severe sepsis or septic shock for the purpose of treating septic shock (grade 1A).
- that corticosteroids not be administered for the treatment of sepsis in the absence of shock. (grade 1D).

# **Adrenal Insufficiency: Current Practice 2012**

**Lawrence S. Kirschner, MD, PhD**  
**Professor of Medicine**

**Division of Endocrinology, Diabetes, and Metabolism**  
**The Ohio State University's Wexner Medical Center**

## **Treatment of Adrenal Insufficiency**

## **Treatment of Adrenal Insufficiency: Glucocorticoid Replacement**

- **Hydrocortisone**
  - Metabolized to cortisol
  - Approx 10-12 mg/m<sup>2</sup> is replacement dose of HC
  - In most people, this is about 20-25 mg/day
    - 5'9", 155 lb patient. BSA = 1.85. Dose = 18-22 mg
    - 6', 300 lb patient. BSA = 2.63. Dose = 26-31 mg
  - Mimic the diurnal variation (2/3 steroid A.M.; 1/3 evening)
    - Evening dose given mid afternoon (e.g., 3pm) unless patient is night owl
  - Can also be given as single AM dose if patient tolerates

## **Glucocorticoid equivalents**

- **Hydrocortisone: 20 mg**
- **Cortisone acetate: 25 mg**
- **Prednisone 4-5 mg**
- **Prednisolone 5 mg**
- **Dexamethasone 0.75-1 mg**
  - Synthetic steroids have longer half life, and may have increased incidence of side effects (e.g., osteoporosis, weight gain, immune suppression)

## **Treatment of Adrenal Insufficiency: Mineralocorticoids**

- Replacement of mineralocorticoid needed if primary adrenal failure
- Florinef is synthetic mineralocorticoid (fludrocortisone)
  - Comes in only 1 size (100 mcg)
  - Most patients need 1 tab/day, but may need to titrate to symptoms or electrolytes
  - In patients on high dose HC (>50 mg/day), enough MC activity so that florinef not usually needed

## **Treatment of Adrenal Insufficiency: Androgens**

- Anecdotal evidence suggests that replacing DHEA(S) may help improve patient well-being, but randomized studies have all been **NEGATIVE**
- Patients may benefit from a trial of DHEA 50 mg.
  - Patient feels better → great!
  - No better → stop.

## **Treatment of Adrenal Insufficiency: Efficacy**

- There is no single lab test that will judge adequacy of replacement, so patient symptomatology important
- ACTH goals:
  - Generally, aim for AM ACTH 50-150 pg/ml [normal 10-50]
  - Lower ACTH values generally indicate overtreatment
- Renin goals:
  - Normalized
  - Note that it may be very difficult to control ACTH levels if patient has significant mineralocorticoid deficient

## **Recovery from Addison's?**

- Patients who fail an ACTH stim should be retested to verify
- Patients with Cushing Syndrome that are cured by surgery will be insufficient until their axis recovers
  - Requirement for steroids post-op is a *good sign*
  - Patients with Cushing syndrome can take 1-2 years to recover
- Patients on chronic steroids for many years may take many years to recover their axis

## **Recovery from critical illness**

- **Patients that are suspected of having adrenal insufficiency should have their steroids weaned once critical illness has resolved**
- **Typically, patients can be weaned to replacement level treatment at the time of discharge**
- **Further evaluation and tapering can then be done in the outpatient setting**

## **Facilitating HPA recovery**

- **Use Hydrocortisone**
  - **Shorter biological half-life means axis can recover while patient on therapy**
  - **If patient tolerates, put on once daily replacement of HC and wait**
    - **Go for lowest dose that patient will tolerate**
  - **Retest by ACTH stim q3-4 months until recovery**
- **Can also use prednisone (low dose or qod dosing) but usually doesn't work as well**