

Environmental Injuries

Colin G. Kaide, MD, FACEP, FAAEM, UHM
Associate Professor of Emergency Medicine
Board-Certified Specialist in Hyperbaric Medicine
Specialist in Wound Care
The Ohio State University Wexner Medical Center



The Most Dangerous Drug Combination...




Testosterone and Alcohol!

The most likely victims...

Photo: Ralf Roletschek

Accidental Hypothermia



Blizzard of



1979

Definition of Hypothermia

- * Subnormal T° when the body is unable to generate sufficient heat to sustain normal functions
- * Core Temperature $< 95^{\circ}\text{F}$ (35°C)

Most Important Temperatures

- * 95°F (35° C) Hyper/Goofy
- * 90°F (32°C) Shivering Stops
- * 80°F (26.5°C) Vfib, Coma
- * 65°F (18°C) Asystole

Thermoregulation

- * The body uses a Poikilothermic shell to maintain a Homeothermic core
- * Maintains core T° w/in 1.8°F(1°C)
- * Hypothalamus
- * Skin

Constant T° 96.8-
100.4° F

Thermoregulation

The 2 most important factors



- * Shivering (10x increase)
- * Initiated by low skin temperature
- * Warming the skin can abolish shivering!
- * Peripheral vasoconstriction
- * Sequesters heat

Only 3 Causes!

- Decreased Heat Production
- Increased Heat Loss
- Impaired Thermoregulation



Predisposing Factors

Decreased Production

- Endocrine problems
 - Thyroid
 - Adrenal Axis
- Malnutrition
- Neuromuscular disease



Predisposing Factors

Increased Loss

- * Radiation
- * Evaporation
- * Conduction*
- * Convection**

***Depends on conducting material**

****Depends on wind velocity**

Predisposing Factors


Impaired Regulation

- * CNS injury
 - * Hypothalamic injuries
- * Peripheral Injury
 - * Atherosclerosis
 - * Neuropathy
- * Interfering Agents

Systemic Responses

CNS

- * $T^{\circ} < 90^{\circ}\text{F}$ (34°C)
 - * Hyperactivity, excitability, recklessness
- * $T^{\circ} < 80^{\circ}\text{F}$ (27°C)
 - * Loss of voluntary motion and reflexes
- * $T^{\circ} < 75^{\circ}\text{F}$ (24°C)
 - * Loss of corneal & oculocephalic reflexes



The patient can look dead!

Systemic Responses


Cardiovascular

- * Above 90°F (32°C): Excitatory
 - * Tachycardia, Hypertension
- * Below 90°F (32°C): Inhibitory response
 - * Bradycardia at level of pacemaker cells
 - * Atrial and ventricular dysrhythmias
 - * Atrial Fib
 - * Vasodilatation




The Infamous Osborn Wave

A form of early repolarization



Systemic Responses

Pulmonary




- * Initially tachypnea is seen
- * $T^{\circ} < 90^{\circ}\text{F}$ (32°C) RR can fall to 5-10

**Minute volume falls
proportional to metabolic rate**

Systemic Responses

Renal




- * Cold diuresis occurs early
- * Central hypervolemia
- * ADH suppression
- * $T^{\circ} < 90^{\circ}\text{F}$ (32°C): Kidney function declines

Systemic Responses

Hematologic

- ❄ Cold coagulopathy can produce significant bleeding
- ❄ Occurs by 2 mechanisms




Systemic Responses

Hematologic

- ❄ Platelet Issues
- ❄ Thrombocytopenia
- ❄ Platelet dysfunction



- ❄ Coagulopathy
- ❄ Dysfunction



Question?

Although bleeding occurs, the measured PT and PTT are usually normal - Why does this happen?

Question?

Do people really take off their clothes when freezing to death?




Image from Ponder High School

Cold Water Immersion- Hmmm?



Image from Chicago Tribune


Cold Water Immersion

- ❄ Heat loss in cold water is 20-30 times that of air!
- ❄ Many additional factors come into play in cold water immersion!

Cold Water Death!

- * Death may occur in only 15 minutes but NOT from hypothermia
- * Cardiac dysrhythmias-“Sudden Disappearance”
- * Breathing abnormalities
 - * Gasp, Hyperventilation
- * Muscular dysfunction

January 13, 1982 Air Florida Flight 90



In water for 1 hour and 45 minutes before rescue!

News Story - Chicago Tribune

Hypothermia Effects

- Video 1: Air Florida Flight 90



"Seconds from Disaster--The crash of Air Florida Flight 90

Video from National Geographic

Jimmy Tontlewicz

"A heroic story of survival"

On Jan. 15, while sledding with his father, Jimmy plunged into the icy waters of Lake Michigan. When rescuers pulled him out, he had been submerged for at least 20 minutes and had no discernible heartbeat, pulse or breathing. In Chicago last week doctors said Jimmy is progressing so well that they hope to send him home this month.



Diver lifts four-year-old Jimmy Tontlewicz out of Lake Michigan

In icy lake for 20 minutes, boy brought back to life



News Story - Chicago Tribune

Management and Rewarming of a Hypothermic Patient

Emergency Medicine Dogma

* You're not dead until you're...

Warm and dead

Unless...

...You're already dead!



Drawn by friend-Kestutis Boyev

Death!




- * Still dead with $T^{\circ} > 32^{\circ}\text{C}$ (90°F)
- * Serum potassium > 10
- * *Documented and verified DNR orders!*

Treatment

Do Primary/Secondary Survey

Evaluate for other treatable conditions



College newspaper: Daily Illini

Rewarming



Most Important Rule

If the core temperature is less than 90°F (32°C) and shivering has stopped, YOU MUST ADD HEAT to the core!



Rewarming is Additive

+ **+** **+** **+** **+** **+** **+**



Rewarming: *Passive*

- ❁ Insulate and allow shivering to raise body temperature
- ❁ Appropriate for mild hypothermia
- ❁ Core Temperature > 32°C/90°F
- ❁ Healthy individuals



Rewarming: *Active*

ADD HEAT!



Rewarming: *Core* *Rewarming*

- ❄ Often *VERY* complicated to perform
- ❄ Supplies heat directly to the core
- ❄ Necessary for most patients with a core temp $< 90^{\circ}$
- ❄ Fast!



Treatment: Core Rewarming

*** Heated Humidified Air**

- * Heat to 45°C/113°F**
- * 2-3°F rise in T°/hr**
- * Indicated for ALL significantly hypothermic patients**



What About heated IVFs?

Is heated IVF an effective rewarming method?

NO!

Why is This?

**70 Kg person is 60% water:
– 42 L of fluid**



**If 42 L of fluid is at 85°F and you add 1
additional L of fluid at 110°F...How much
difference does it make?**

Only 0.3°C/0.6°F per Liter

Treatment: Core Rewarming


- * Heated irrigation of body cavities
 - * Abdominal Irrigation (“TPL”)
 - * ~3°F/hr
 - * Thoracic Irrigation
 - * Very effective (up to 10°F/hr)
 - * Ant/post chest tubes

Treatment: Therapeutic Peritoneal Lavage



Treatment: Thoracic Irrigation

- Video 2: Thoracic Irrigation



Treatment: Rewarming

- * Extracorporeal Blood Rewarming
- * Fem-Fem Bypass
- * CAVR-Level 1 Infuser
- * V V R
- * Dialysis with a heat exchanger




Microwaves?




Treatment: Sinus Bradycardia

- * Physiologically normal at $T < 93^{\circ}\text{F}/34^{\circ}\text{C}$
- * Don't treat it—Self-limited



Treatment: Atrial Fibrillation

- * Occurs Commonly at $T < 86^{\circ}\text{F}$ (30°C)
- * The rate is **SLOW!**
- * Resolves with treatment of Hypothermia



Treatment: Ventricular Fib

- * Occurs $< 28^{\circ}\text{C}/83.5^{\circ}\text{F}$
- * Lidocaine is ineffective
- * Rewarm and defibrillate every few degrees





Treatment: Asystole

- ❄ Occurs physiologically at $T < 65^{\circ}\text{F}/18^{\circ}\text{C}$
- ❄ May occur spontaneously
- ❄ Only responds to rewarming



Other Treatments

- ❄ CPR
- ❄ Only when no detectable pulse
- ❄ Pressor agents
- ❄ Caution with cardiac stimulation

Heat-Related Illness




Statistics

- About 500 die each year in the U.S.
- Hard to know exact number because it's often under-reported
- August 2003: at least 35,000 died in Europe

Chicago, August, 1995



Photo: Daniel Schwen



All tolled, 760 people, mostly the elderly and poor died that summer

Pathogenesis of Heat Illness

- Exogenous heat gain
- Endogenous heat production
- Decreased dissipation

Pathogenesis:

Exogenous Gain

- Environmental temperature
 - Sun, workplace, home, sauna



Wet Bulb Globe Temperature! A Weighted Average...



- 10%: Dry, shaded thermometer
- 70%: Wet thermometer
- 20%: Unshaded black globe

Pathogenesis: Endogenous Production

- **Basal metabolism: 50-60 kcal/hr/m²**
- **1°C/hr increase in T° if we had no mechanism for dissipation!**

20x Increase in heat production is seen during exercise!

Pathogenesis: Endogenous Production

- **What are some other causes...**
 - **Hyperthyroidism**
 - **Neuroleptic Malignant Syndrome**
 - **Malignant Hyperthermia**
 - **Cocaine, Amphetamines, MDMA, LSD**
 - **Fever**

Pathogenesis: Decreased Dissipation

Yeah...But
it's a dry
heat!



- Dehydration is the *most* significant factor affecting the ability to dissipate heat!

Hullraisere on Panoramio.com

Decreased Dissipation: Dehydration

- Limits Sweating
 - Volume overrides heat dissipation
- Impairs CV function
- Insensible water loss
 - 1.5L/day (2% BW)
 - Exercise: 1-2 L/hr
- Maximum gastric emptying
 - 1-1.5 L/Hr

QUESTION?

How much of the lost fluid does thirst alone replace?

Only about 2/3 of the needed fluids



Spectrum of Illness

**Heat
Edema**

**Heat
Cramps**

**Heat
Syncope**

**Prickly
Heat**

**Heat
Exhaustion**

**Heat
Stroke**

Heat Exhaustion

- Flulike symptoms – malaise, headache, weakness, nausea, anorexia, vomiting
- Tachycardia, orthostatic hypotension
- Sweating is generally present
- Temperature is $< 40^{\circ}\text{C}$ (104°F)
- Mental status and neurologic exam are normal

Heat Exhaustion

- Cool shaded environment
- Oral rehydration if capable but may need IVF due to large amounts of volume lost as sweat
- Cooling is not necessary but it can make the patient feel better

**What is the Most Important Thing to
Tell a Discharged Heat Exhaustion
Patient???**



Heat Stroke

- **Catastrophic, life-threatening emergency**
 - **Failure of thermoregulatory mechanisms**
 - **Multisystem tissue/organ damage**
- **Damage is a function of T° max and duration of T° elevation**



Heat Stroke

- **Temperature $> 40.5^{\circ}\text{C}$ (105°F)**
- **MENTAL STATUS CHANGES:**
Hallmark is severe CNS dysfunction
 - **Confusion**
 - **Delirium**
 - **Seizures**
 - **Coma**



Multi-Organ Dysfunction


- Encephalopathy
- Rhabdomyolysis
- Acute renal failure
- ARDS
- Myocardial/hepatocellular/pancreatic
- Intestinal ischemia/infarction
- Bleeding complications – DIC



www.gearfuse.com

Heat Stroke: *Area of Confusion*

- Can the temperature be less than 105°F and still be heat stroke??



Heat Stroke: *Area of Confusion*

- Anhydrosis (sweat cessation)
 - Sweat gland fatigue
 - Dehydration
- Sweating can persist to $T^{\circ} > 42^{\circ}\text{C}$ (108°F)

Classic (Epidemic) Heat Stroke

- Excess heat gain, impaired loss
- Occurs during heat waves
- Elderly, very young, poor, debilitated
- +/- inciting medications
- *Sweating is less likely*



politicalhumor.com

Exertional Heat Stroke

- XS heat production, overwhelmed loss mechanisms
- Young, healthy, athletes, military, etc.
- Worse systemic involvement
- Rhabdo, ARF, coagulopathy, hypoglycemia
- *More likely to still be sweating*

How long can it take a runner in 100% humidity at 85°F to develop heat stroke?

Vikings football player dies of heat stroke
August 2, 2001 Posted: 6:26 AM EDT (1026 GMT)

Korey Stringer died early Wednesday of heat stroke



Image from CNN

IRIE, Minnesota (CNN) -- Pro Bowl lineman Korey Stringer of the Minnesota Vikings died of heat stroke early Wednesday, the

4, 335-pound Stringer, 27, died at St. Joseph's Hospital -- Mayo Health Mankato, where the team holds its practices.

The death came as dangerously hot weather continued to pose a problem for the central United States. Temperatures were expected to reach as high as 100 degrees in Iowa and Illinois on Wednesday.

Stringer began exhibiting signs of heat stroke, including weakness and rapid breathing, after a morning practice session Tuesday.

Heat Stroke Claims Local Football Player
August 12, 2005

OKLAHOMA CITY -- Medical examiners said that an autopsy done on Douglass High School football player Chris Stewart Friday determined that the 17-year-old died from heat stroke.

Stewart collapsed at a Tuesday practice, in 95-degree heat. He was taken to the hospital with elevated blood pressure and body temperature and with some brain swelling. Prior to the collapse, Stewart was projected to be a starter on the Trojans offensive line, and his teammates said they were unaware of any pre-existing medical conditions Stewart may have had.

Stewart was also an honor student. Tests are pending, but expected to be completed Wednesday at 11 a.m.



Image from CNN

Treatment: Cooling

- Evaporative cooling (Khogali method)
- 15°C (59°F) mist + Fan 45°C (113°F)
 - 0.06°C (0.1°F)/min



The Evaporative Method



TX: Cooling: Ice/Cold Water Immersion



- 0.13-0.16°C decrease/min (0.23-0.28°F)

Aggressive Resuscitation



Treatment

- ABCs
- IVF – treat volume depletion
- Avoid shivering
 - Benzodiazepines for seizures/shivering
- Dantrolene is ineffective
- Monitor for complications and treat


Good Prognosis

- **Recovery of central nervous system function during cooling**
- **Expected in the majority of patients who receive prompt and aggressive treatment**

Poor Prognosis

- **Coagulopathy with liver hepatocyte damage**
 - **AST > 1000 U/L**
- **Lactic acidosis in classic form**
- **Rectal temperature > 108° F**
- **Prolonged coma**

Don't Forget Your Furry Friends!



photowebster



Dysbarism


- **All the pathologic changes caused by altered environmental pressure**
 - **Altitude-related event**
 - **Underwater diving accident**
 - **Blast injury that produces an overpressure effect**

Types

- **Barotrauma – Expansion of trapped gases**
- **Decompression sickness – Gas bubble disease**

Pressure


Top of Atmosphere



- 14.7 psi
- 33 ft seawater
- 10 m seawater
- 1 atm
- 760 mmHg
- 760 Torr

Pressure

- At 33 ft of seawater




Flying

- Most commercial aircraft are pressurized to 8000 ft
- 0.73 ATA
- FiO_2 21% but functionally less molecules of oxygen per breath ~ 16% FiO_2



Gas Laws: Boyle's Law


- “The volume of a gas is inversely proportional to the pressure exerted upon it”



Consequences of Pressure



Middle Ear Squeeze- *Barotitis media*

- Most common diving-related barotrauma
- Failure to equalize
- Too rapid descent or infection/inflammation
- TM is pushed inward and can rupture



Other Barotrauma

- Barosinusitis
- Barodontalgia
- Alternobaric vertigo
- Face mask squeeze



Scuba Rule # 1

**Never Hold
Your Breath!**




Breath-holding Kills

Blowing Bubbles



Exogenous Entry of Air




Pulmonary Over-Pressurization

- **Can get:**
 - **Pneumothorax, pneumomediastinum, SQ emphysema, rupture into pulmonary vein causing air embolism**
- **Simple pneumothorax may progress to tension on further ascent**

Air-Gas Embolism (AGE)


- **Bubbles enter the pulmonary venous circulation from ruptured alveoli**
- **Usually develops right after diver surfaces**
 - **Sudden LOC = Air embolus until proven otherwise**
- **Cardiac**
 - **Ischemia—dysrhythmias, cardiac arrest**
- **Neurologic**
 - **LOC, confusion, stroke-like sx**

Cerebral Air-Gas Embolism—CAGE



Hyperbaric Oxygen and Bubble Reduction

- As pressure increases, the bubble size decreases and O_2 replaces the inert gas in the bubble (N_2), which promotes diffusion




Air Embolism

- Recompression in hyperbaric chamber
- Transport supine
- 100% oxygen, intubate PRN
- IVF
- Aspirin for antiplatelet activity if not bleeding
- Lidocaine

Decompression Sickness (DCS)

- Henry's Law: "The amount of gas (O_2 and N_2) dissolved in a liquid (blood plasma) is proportional to its partial pressure"



DCS


- The longer and deeper the dive, the more nitrogen gas will be accumulated in the body

DCS



DCS

Slow ascent allows for compensation



Type I DCS



- Periarticular joint pain is the most common symptom of DCS
- Dull, deep ache that is mild then more intense
- Palpable tenderness
- "The Bends"



Public domain in the United States

Type I DCS

- **Cutaneous**
 - Pruritus
 - Cutis marmorata
 - Hyperemia
 - Orange peel
- Lymphedema
- Fatigue, especially if severe



Type II DCS = 10-15%

- Nervous system
- Pulmonary system (< 2%)

Neurologic DCS

- **Spinal cord is most common site**
- **Lower thoracic and lumbar regions**
 - **Low back pain**
 - **“heaviness” in legs**
 - **Paresthesias**
 - **Possible bladder or anal sphincter dysfunction**
- **Brain – variety of symptoms and difficult to distinguish from AGE**
 - **Scotomata, headache, confusion, dysphasia**

Pulmonary DCS

- **“The Chokes”**
- **May begin immediately after dive but often takes up to 12 hours to develop**
- **Triad – shortness of breath, cough, and substernal chest pain or chest tightness**
- **Cyanosis, tachypnea, and tachycardia**

DCS Treatment


- ABCs
- 100% oxygen
- IVF
- *Recompression therapy*
- Divers Alert Network (DAN): 919-684-8111
- 75-85% have good results when recognition and treatment are prompt

Delivery of Therapy

- Monoplace Chambers
 - Single patient



Multiplace Chamber



Multiplace Chambers

