

Gastrointestinal Bleeding

Bennie Ray Upchurch III, MD, FACP, FASGE
Clinical Associate Professor of Medicine
Division of Gastroenterology, Hepatology & Nutrition
The Ohio State University Wexner Medical Center

Disclosures

- **Nothing to disclose**

Objectives

- **Develop a prioritized care plan for patients admitted with upper gastrointestinal bleeding (UGIB).**
- **Accurately assess, triage and resuscitate the UGIB patient.**
- **Recognize common causes of UGIB and the approach in management.**
- **Understand and facilitate a multidisciplinary approach to management of the UGIB patient.**

Acute Upper Gastrointestinal Bleeding

- **Annually ~ 300,000 hospitalizations and ~ 20,000 deaths in US***
- **Common cause for ICU admission and potentially lethal medical emergency**
- **Overall incidence: 50-100/100,000 pts/yr**
- **Incidence of UGIB : LGI bleeding ~ 5:1**
- **More common in elderly – esp. men > 70 yrs, who comprise ~ 30 % of all pts with UGIB**
- **Mortality rates over past 40-50 yrs had been stable at ~ 7-10%. More recently improved to ~ 2.4-5%**

*Acute non-variceal UGIB

Etiology of UGIB

- **Non Variceal: 86%**
 - **Ulcerations: about 50%**
 - **Mallory-Weiss Tear: 4-8%**
 - **Erosive esophagitis: 1-13%**
 - **Neoplasia: 2-7%**
 - **Vascular ectasia: 0-6%**
- **Variceal: 14%**

Initial Assessment of Severe UGIB

- 1. Resuscitation and stabilization**
- 2. Assessment of severity and location of bleeding**
- 3. Preparation for emergent upper endoscopy**
- 4. Role of endoscopist**
 - **Localization and identification of the bleeding site**
 - **Control of active bleeding or high risk lesions**
 - **Stratification of the risk for rebleeding**
 - **Minimization of treatment-related complications**
 - **Treatment of persistent or recurrent bleeding**

Initial Evaluation - History

- **Age: Elderly (ischemia, cancer, diverticula)**
Young (ulcers, esophagitis, varices)
- **Prior GI Bleeding**
- **Previous gastrointestinal disease**
- **Previous GI surgery**
- **Underlying Medical Disorders (liver disease, CKD)**
- **Meds : NSAIDS - ASA/Anticoagulant use**
 - **Symptoms: Abdominal pain, fever, wt loss, anorexia, epistaxis, hematuria etc**

Physical examination

- **Hemodynamics with a thorough cardiopulmonary exam**
- **Skin (spider angiomas, purpura, cutaneous telangiectasias /pigmentation)**
- **Abdomen (ascites, tenderness, masses)**
- **Digital Rectal exam**

Initial Patient Care/Management

- **Appropriate IV access: 2 large bore I.V. catheters**
- **IV fluids (NS/LR) and/or blood product resuscitation.**
(Target HCT 30% in elderly/ 25-30% in young adults and pts with in Portal HTN)
- **Continuous cardio-pulm monitoring for those with coronary risk factors with supplemental O2.**
- **Frequent vital sign / urine output monitoring.**
- **Consider intubation in those with altered mental status or brisk bleeding.**

Labs and Studies

- **Complete blood count, electrolytes (BUN >> Cr)
Albumin for risk scoring. Consider Iron panel.**
- **Coagulation Panel: PT/INR**
- **Type + screen or type + cross-match blood**
- **EKG for patients > 50 yrs or risk factors for heart disease.**
- **Abdominal radiographs usually not indicated.**

RBC transfusion

- **Restrictive strategies**
 - Transfusion may disrupt splanchnic vasoconstriction, increase splanchnic BP, impair clot formation
 - Threshold of $\leq 7\text{g/dL}$ assoc with lower mortality in ill patients, higher 6 wk survival, and lower rebleeding rate
 - Physician's judgment in active bleeding & with comorbidities

Transfusion Strategies for Acute Upper Gastrointestinal Bleeding
C Villanueva, et al. N Engl J Med 2013; 368:11-21

Naso Gastric Lavage (NGL)

- 32% positive predictive, 85% negative predictive value
- Positive NGL does not provide etiology
- A non bloody aspirate in ~ 25% of UGIB
- A bile aspirate does not R/O UGIB
- Minimal evidence that NGL affects outcome (risk scoring)

Aljebreen AM, Fallone CA, Barkun AN. Nasogastric aspirate predicts high- risk endoscopic lesions in patients with acute upper-GI bleeding. Gastro- intest Endosc 2004;59:172-8.

Huang ES, Karsan S, Kanwa IF, et al. Impact of Nasogastric lavage on outcomes in acute GI bleeding. Gastrointest Endosc 2011;74:971-80.

Risk Scoring

Rockall

- Pre and post-endoscopy values
- Predicts high or low risk for rebleeding, mortality
- Prospectively pre-endoscopic score less reliable for low risk.

Glasgow Blatchford

- Pre-endoscopy values
- Predicts need for interventions
 - Endoscopic
 - Surgery
 - transfusions

Causes of upper GI bleeding in hospitalized patients

- Gastroduodenal erosions 44%
- No source found 23%
- Esophagitis 22%
- Other 15%
- Gastric Ulcer 8%
- Duodenal Ulcer 2%
- Multiple findings

Uncommon Sources of GI Bleeding


- **Hemosuccus Pancreaticus**
- **Hemobilia**
- **Dieulafoy lesion**
- **Vascular Ectasias**
- **Aorto-enteric fistulae**
- **Neoplasms : Benign > Malignant**


Predictors of Mortality

- **Increasing age - Age > 70 yrs**
- **Concurrent active major organ disease**
- **Preexisting hospitalization (mortality rate ~ 34%)**
- **Passing frequent frank blood. Esp if - Shock or Orthostatic hypotension**
- **Requiring emergency surgery for GIB**
- **Active bleeding / Transfusion requirement**
 - **4 or more red cell units in the first 24 hours**
 - **2 or more units for rebleeding event**
 - **6-8 units total**

Forrest Classification System with Respective Prognosis

Forrest Classification	Rebleeding Incidence	Surgical Requirement	Incidence of Death
Type I: <u>Active</u> Bleed Ia: Spurting Bleed Ib: Oozing Bleed	55-100%	35%	11%
Type II: <u>Recent</u> Bleed IIa: Non-Bleeding Visible Vessel (NBVV) IIb: Adherent Clot	40-50% 20-30%	34% 10%	11% 7%
Type III: Lesion <u>without</u> Bleeding Flat Spot Clean Base	10% 5%	6% 0.5%	3% 2%





Management of UGI Bleeding

- High dose PPI Rx
- Endoscopic Modalities
 - Injection Rx
 - Thermal device
 - Mechanical

Endoscopic Methods of Hemostasis of UGIB

- **Thermally active**
 - Heater probe
 - APC
- **Injectable therapies**
 - Epinephrine
 - Glue
- **Mechanical**
 - Endoscopic clips
 - Band ligation
- Combination Rx

Prognostic Features of GastroDuodenal Ulcers

- **Posterior duodenal wall or lesser gastric curvature**
- **Ulcer size > 1 cm is associated with increased re-bleeding and mortality**
- **Endoscopic hemostasis is less successful in ulcers > 2 cm in size**
- **Greatest re-bleeding risk from ulcers is within first 72 hours**

H. pylori (H.P) and PUD

- **ASGE and ACG guidelines suggest that all pts with PUD should be checked and treated for H. pylori. (Class A rec)**
- **Some studies suggest rebleeding less with H.P Rx than PPI.**
- **NSAID user infected with H.P has ~ two-fold risk of ulcer bleeding.**
False negatives/false positives
- **Theoretically, alkaline milieu in UGIB (or PPI use) results in proximal migration of H.P**
- **Serologic testing is unreliable for active infection or proving eradication**

PUD and NSAIDS / ASA

- **Mechanism : Reduced production of cyclooxygenase –generated cytoprotective PG, platelet dysfunction**
- **Risk of Bleeding : gastric ulcers > duodenal ulcers**
- **RR of NSAIDS is 4-7 compared to ASA (2.5) and COX 2 inhibitors (1.5). Relative risk varies with individual NSAIDS ex : piroxicam > ibuprofen etc**
- **Risk of bleeding is dose dependant**
- **Multiple cofactors contribute to risk (eg. age > 75 yrs, h/o CAD, prior GIB, H. pylori, steroids, bisphosphonates & ETOH etc)**

Management while on ASA

- **Consensus recommendation-Short term hold for 5 days**
- **Restart as soon as the risk for cardiovascular complication outweighs risk for bleeding**
- **3 fold increase in major cardiac events within 7-30 days**
- **Gastroprotection with PPI (CONGENT)**
 - **Continue PPI as long as on ASA/DAPT**
 - **PPI with Plavix if history of PUD**

Resumption of Low Dose ASA (81 mg) After Bleeding Ulcer

- **Sung et al . Gastro 2006;130 (suppl 2): A 44**
- **8 week DB RCT after Endoscopic Therapy. IV PPI X 3 days followed by oral PPI Rx.**
- **Rebleeding at 1 month; 11% of pts on placebo (N=55) rebled c/to 18% of pts on ASA (N=58).**
 - **P = 0.25**
- **Mortality at 2 months: There was a 14% mortality in placebo group (n=55) vs 2% in pts on ASA (N=58).**
 - **P = 0.012.**

Medical Therapy for Bleeding PUD

- Acidic pH retards clotting, enhances clot dissolution.
- PPI : Clearly superior to H2RA to keep gastric pH > 6.0
- Pre endoscopic PPI compared with H2RA showed no evidence of reduced rebleeding, need for surgery, or mortality
- Pre endoscopic PPI downstages high risk lesions to low risk
- Post endoscopic high dose intravenous PPI therapy (80mg bolus dose intravenously followed by 8mg/h infusion over 72 hours) (Class A rec)
- All patients should be discharged on a single daily oral PPI dose. Caveat- GERD

Prokinetics

- Erythromycin 250mg IV 30-90 min before EGD
 - significantly increases quality of mucosal visibility.
 - (Class A rec)
 - reduces need for re look endoscopy
 - Consider EKG
- Reglan 10mg IV 30 min prior (option)
 - Caution regarding tardive dyskinesia/EPS
- Evidence based > NG lavage

Rebleeding after Endoscopic Therapy

- ~ 20% of pts with active UGIB rebleed.
- A 'second-look' endoscopy demonstrated benefit in only those cases with active re-bleeding.
- 3386 patients with bleeding peptic ulcers
 - Initial therapy 98.6% successful
 - Rebleeding 8.2%
 - Predictors of rebleeding:

	OR
•Hypotension	2.2
•Anemia <10 gm/dl	1.9
•Active bleeding / fresh blood	1.7 / 2.2
• \geq 2 cm ulcer	1.8

Wong et al Gut 2002

Early (2-24 hrs) vs. Delayed Endoscopy for UGIB

- Lower costs
- Early discharge of low risk patients.
- Location of admission (ICU vs. ward)
- Significant benefit of endoscopic therapy in high risk pts has not been documented in RCTs.
- Major clinical outcome parameters such as rebleeding rate, mortality and the need for an emergency operation have no bearing with timing of endoscopy.

J Sung, AGA Perspectives Vol 5, Dec 2009

Indications for Angiography in UGIB

- **Consensus statement from American College of Radiology:**
- **Endoscopy is the best dx and therapeutic procedure.**
- **Surgery and Transcatheter arteriography /intervention (angiography) are equally effective following failed EGD.**
- **Angiography considered in cases with high operative risk**
 - **less successful in pts with impaired coagulation**
 - **best technique for UGIB into the biliary tree or pancreatic duct.**

Angiographic Therapy

- **Overall - is rarely required in pts with bleeding ulcer.**
- **Bleeding should be > 0.5 ml/hr.**
- **Selective Intra-arterial vasopression – not used now.**
Risks: Brady-arrhythmias, ischemia, etc
- **Selective occlusion of bleeding arteries with gelfoam, beads, tissues adhesives and coils etc are used.**
- **Rebleeding is common, and complications such as ischemia, infarction, perforation and abscess etc are prominent.**

Surgical Therapy for UGIB- When ?

- Role is controversial.
- Is usually considered in high risk cases when;
 - 1) HD instability even after > 3 units PRBC transfusions
 - 2) TWO unsuccessful EGDs/attempts at hemostasis
 - 3) Shock with recurrent hemorrhage
 - 4) Continuous bleeding with transfusion requirements of > 3 units PRBC / day.

Surgical Therapy

- Typically pts are severely ill and mortality is ~ 25 %
- Primary objective is not to cure ulcer disease but stop hemorrhage. Acid-reducing procedures may be added.
- A large RCT trial of 92 pts – demonstrated that after initial failure of Endo Tx – an endoscopic re-treatment reduced the need for surgery without increasing mortality and had fewer complications than surgery.
- No data from current endoscopic era supports early surgery except - A-E fistula, bleeding benign tumors and severe GAVE

Stress Related Mucosal Disease

- Incidence decreasing since 90's with improved ICU care.
- In ventilated pts with respiratory failure (OR - 15.6) and/or coagulopathy (OR - 4.3).
- Prophylaxis with H2RA > sucralfate (GIB 1.7% vs. 3.8% , P=0.02).
- H2RAs may be limited by tolerance.
- PPIs may have possible interactions with Plavix and potential increased risk of C. difficile infection.
- No pharmacotherapy shown to be beneficial once bleeding.
- Endoscopic therapy should be attempted.

UGIB after AMI

- **Not uncommon**
 - 1-3%
 - Multifactorial
 - Medications
 - Underlying low flow state

UGIB after AMI

- **Predictors of UGIB**
 - Older age
 - Hemodynamic compromise
 - Severe myocardial ischemia
 - Use of thienopyridines before event
 - +/- Integrillin
- **Substantially increased mortality**
 - Especially if PCI done for NSTEMI/ACS
- **PPI use provides substantial risk reduction**

UGIB after AMI

- **EGD/Colonoscopy is relatively safe in patients with UGIB and AMI**
 - NG Tube is also safe
- **Diagnostic yield is approximately 80%**
- **UGIB followed by AMI *may* be worse than AMI followed by UGIB**
- **Cardiac status may play a less prominent role in complications**
- **Prospective data are sorely needed**

Therapies For Long-Term Prevention of Ulcer Hemorrhage

- **Medical therapies**
 - **Acid suppression**
 - **Prostaglandin analogs**
 - **Mucosal protectants**
- **Helicobacter Pylori eradication**
- **NSAID discontinuation**
- **Smoking cessation**

Conclusions:

- **Medical stabilization**
- **Signs and symptoms help to localize**
- **Direct the investigation**
- **Maximize pharmacotherapy**
- **Alter risk factors**