



## Advances in Liver Disease

**Vivek Mendiratta, MD**  
*Assistant Professor - Clinical  
Transplant Hepatology  
The Ohio State University Wexner Medical Center*

**MedNet21**  
Center for Continuing Medical Education

 **THE OHIO STATE UNIVERSITY**  
WEXNER MEDICAL CENTER

## Disclosures

- No financial disclosures

## Goals and Objectives

- Updates in steatotic liver disease
- Novel treatments for hepatocellular carcinoma
- Autoimmune liver disease, a changing of the guard?



## Case Presentation

52 y/o male with PMHx significant for Type II DM, Hypertension, Obesity (BMI: 38), and Hyperlipidemia presents to clinic for elevated LFTs.

- Patient is new to clinic from another state
- Has two tattoos, professionally done
- Drinks approximately three to four drinks per week
- No history of blood transfusions
- Physical exam notable for obesity

- Liver profile:
  - AST: 34
  - ALT: 55
  - Alk Phos: 78
  - Total Bilirubin: 0.7
  - Direct Bilirubin: 0.1
  - Total Protein: 7.9
  - Albumin: 4.1

- Hepatitis panel negative
- PETH negative
- A1AT negative
- Iron studies normal
- US Abdomen showed hepatic steatosis without overt nodularity

## Steatotic Liver Disease: Scope of the Problem

Second most common cause of HCC and Liver Transplantation in the United States


Disease awareness remains limited

Early detection is critical to preventing fibrosis progression

## Nomenclature Change

- Metabolic dysfunction-associated steatotic liver disease (MASLD) replaces non-alcoholic fatty liver disease (NAFLD)
- Metabolic dysfunction-associated steatohepatitis (MASH) replaces non-alcoholic steatohepatitis (NASH)
- Metabolic and alcohol-related liver disease (MetALD)
  - SLD for people with MASLD who drink more than 140g/week ( $\geq 10$  drinks/week) for females or 210g/week ( $\geq 14$  drinks/week) for males

# MASLD Spectrum



# Global Prevalence






Image source: <https://communityliveralliance.org/liver-health-facts/>

Younossi, Hepatology, Sept 2018


# Increasing Prevalence of MASH and Advanced Fibrosis



Chris Estes, Homie Razavi, Rohit Loomba, et al.. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. Hepatology 67(1):p 123-133, January 2018. | DOI: 10.1002/hep.29466 CC BY-NC 4.0

**MASLD Patients in the US: 100 million**

**Hepatologists: 7500**



## Screening Recommendations

### AASLD 2022

- Type II DM
- Medically complicated obesity
- Hepatic Steatosis with moderate alcohol consumption
- First-degree relatives of a patient with cirrhosis due to NAFLD/NASH

### AACE 2022

- Pre-diabetes or Type II DM
- Obesity and/or  $\geq 2$  cardiometabolic risk factors
- Hepatic steatosis (on imaging) or increase in AST/ALT

### EASL 2016

- Obesity, Type II DM, or Metabolic syndrome
- Persistently abnormal liver enzymes

# Clinical Assessment Tools

## Simple Calculators

- FIB-4
- NAFLD Fibrosis Score

## Serum Tests

- Fibrosure
- Enhanced Liver Fibrosis (ELF)


## Imaging

- Transient elastography
- MR elastography


## Histologic Assessment

- Liver Biopsy

# Simple Calculators




# Imaging Techniques: Fibroscan



MEASURES  
VELOCITY OF  
SOUND  
WAVES  
PASSING  
THROUGH THE  
LIVER




TAKES 5-10  
MINUTES



GIVES  
ASSESSMENT  
OF STEATOSIS  
(CAP SCORE)  
AND FIBROSIS

## Algorithm



Chris Estes, Homie Razavi, Rohit Loomba, et al. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* 67(1):p 123-133, January 2018. | DOI: 10.1002/hep.29466 CC BY-NC 4.0





# Treatment: Lifestyle Modification



- 3% weight loss improves steatosis  
10% weight loss improves fibrosis
- 150-200 minutes of exercise per week
- No alcohol intake if  $\geq$ F2 Fibrosis
- Drink 2+ cups of coffee per day

# Weight Loss Pyramid





## A Placebo-Controlled Trial of Subcutaneous Semaglutide in Nonalcoholic Steatohepatitis


Philip N. Newsome, M.B., Ch.B., Ph.D., Kristine Buchholtz, M.D., Ph.D., Kenneth Cusi, M.D., Martin Linder, M.Sc., Takeshi Okanoue, M.D., Ph.D., Vlad Ratziu, M.D., Ph.D., Arun J. Sanyal, M.D., Anne-Sophie Sejling, M.D., Ph.D., and Stephen A. Harrison, M.D. for the NN9931-4296 Investigators\*

- 72-week, double-blind trial involving patients with biopsy-confirmed MASH (F1, F2, or F3 fibrosis)
- 320 patients were randomly assigned to receive semaglutide at a dose of 0.1 mg, 0.2 mg, 0.4 mg or to receive placebo
- MASH resolution occurred in 40% in the 0.1-mg group, 36% in the 0.2-mg group, 59% in the 0.4-mg group, and 17% in the placebo group ( $P < 0.001$  for semaglutide 0.4 mg vs. placebo)
- Trend towards improvement in fibrosis stage occurred in 43% of the patients in the 0.4-mg group and in 33% of the patients in the placebo group ( $P = 0.48$ )

Newsom NEJM 2020

# Resmetirom

- Drug that acts as an agonist on thyroid hormone receptor- $\beta$ 
  - THR- $\beta$  pathway is active primarily in the liver
  - Regulates de novo lipogenesis
  - Reduces LDL and improves metabolic control
- Originally developed to treat dyslipidemia
- Patients with hypothyroidism have higher rates of MASLD



## A Phase 3, Randomized, Controlled Trial of Resmetirom in NASH with Liver Fibrosis


S.A. Harrison, P. Bedossa, C.D. Guy, J.M. Schattenberg, R. Loomba, R. Taub, D. Labriola, S.E. Moussa, G.W. Neff, M.E. Rinella, Q.M. Anstee, M.F. Abdelmalek, Z. Younossi, S.J. Baum, S. Francque, M.R. Charlton, P.N. Newsome, N. Lanthier, I. Schiefke, A. Mangia, J.M. Pericàs, R. Patil, A.J. Sanyal, M. Nouredin, M.B. Bansal, N. Alkhouri, L. Castera, M. Rudraraju, and V. Ratzu, for the MAESTRO-NASH Investigators\*

- 52-week, double-blind trial involving patients with biopsy-confirmed MASH (F1, F2, or F3 fibrosis)
- 966 patients randomly assigned in a 1:1:1 ratio to receive once-daily resmetirom at a dose of 80 mg or 100 mg or placebo
- MASH resolution achieved in 25.9% in the 80-mg resmetirom group, 29.9% in the 100-mg resmetirom group, and 9.7% in the placebo group ( $P < 0.001$  for both comparisons with placebo)
- Fibrosis improvement by at least one stage achieved in 24.2% in the 80-mg resmetirom group, 25.9% in the 100-mg resmetirom group, and 14.2% in the placebo group ( $P < 0.001$  for both comparisons with placebo)
- Change in LDL from baseline to week 24 was -13.6% in the 80-mg resmetirom group, -16.3% in the 100-mg resmetirom group, and 0.1% in the placebo group ( $P < 0.001$  for both comparisons with placebo)
- Incidence of serious adverse events was similar across trial groups: 10.9% in the 80-mg resmetirom group, 12.7% in the 100-mg resmetirom group, and 11.5% in the placebo group

Harrison, NEJM, 2024

# How should we use it?

- Consider it in patients with F2 or F3 Fibrosis
- Has not yet been studied in patients with compensated cirrhosis (F4 fibrosis) though studies ongoing
- Dosing: 80mg per day if <100kg, 100mg per day if >100kg
- Does not replace lifestyle modifications and should be used in conjunction to nutrition/exercise counseling
- Consider combination therapy with GLP-1 in diabetic or obese patients
- How do we know that it is working?
- How long does the patient need to be on this?



# Case Presentation

- 52 y/o male with MASLD, T2DM, Hypertension, Obesity (BMI: 38) and Hyperlipidemia who presented with elevated LFT's and hepatic steatosis on imaging. What test should we order next?
- Fibroscan showing F2 Fibrosis and S3 Steatosis
- What type of treatment options should we consider in this patient?
- Patient presents years later with abdominal pain. He undergoes an MRI showing a cirrhotic appearing liver, as well as a 4.2cm right hepatic lobe LR-5 observation with tumor thrombus. His AFP is 450 ng/ml.



# What's New?



PRACTICE GUIDANCE

## AASLD Practice Guidance on prevention, diagnosis, and treatment of hepatocellular carcinoma

Singal, Amit G.; Llovet, Josep M.<sup>2,3,4</sup>; Yarchoan, Mark<sup>5</sup>; Mehta, Neil<sup>6</sup>; Heimbach, Julie K.<sup>7</sup>; Dawson, Laura A.<sup>8</sup>; Jou, Janice H.<sup>9</sup>; Kulik, Laura M.<sup>10</sup>; Agopian, Vatche G.<sup>11</sup>; Marrero, Jorge A.<sup>12</sup>; Mendiratta-Lala, Mishal<sup>13</sup>; Brown, Daniel B.<sup>14</sup>; Rilling, William S.<sup>15</sup>; Goyal, Lipika<sup>16</sup>; Wei, Alice C.<sup>17</sup>; Taddei, Tamar H.<sup>18,19</sup>

Author Information©

Hepatology 78(6):p 1922-1965, December 2023. | DOI: 10.1097/HEP.0000000000000466

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

## Hepatocellular Carcinoma

Version 2.2023 — September 14, 2023



# Hepatocellular Carcinoma: Why is it important?



Liver cancer is the 2<sup>nd</sup> most frequent cause of cancer-related deaths

Incidence rates estimated to exceed 1 million by 2025

Screen with US and AFP every six months

2-5% per year risk of HCC in cirrhotic patients

# Management



## IMbrave150: Atezolizumab plus Bevacizumab for Unresectable HCC

- 501 patients with unresectable HCC were randomly assigned in a 2:1 ratio to receive either atezolizumab plus bevacizumab or sorafenib
- Overall survival at 12 months was 67.2% (95% CI, 61.3 to 73.1) with atezolizumab–bevacizumab and 54.6% (95% CI, 45.2 to 64.0) with sorafenib
- Median progression-free survival was 6.8 months (95% CI, 5.7 to 8.3) and 4.3 months (95% CI, 4.0 to 5.6) in the respective groups

Cheng, Journal of Hepatology, 2022  
Finnm, NEJM, 2020

## Himalaya: Tremelimumab plus Durvalumab in Unresectable HCC



Tremelimumab/Durvalumab (n=393) vs. Durvalumab (n=389) vs. Sorafenib (n=389)



OR for STRIDE was 20.1%, 17.0% with Durvalumab alone, 5.1% for Sorafenib




Median Survival: 16.43 months with STRIDE, 16.5 months with Durvalumab, and 13.7 months with Sorafenib




Time to QOL deterioration: 7.5 months for STRIDE, 7.4 months for Durvalumab, 5.7 months for Sorafenib

About-Alfa, July 2022, NEJM

# Transarterial Radioembolization




- Microspheres embedded with Yttrium-90 (Y-90)
- Intra-arterial delivery
- High radiation doses to the target tumor



- Performs better than cTACE
- Longer TTP (26 vs. 6.8 months)
- Improved pathologic necrosis (87% vs. 74%)
- Better side effect profile

# Combination Therapies: Is more better?





## Emerald 1: TACE+Durva+Bev

- 616 patients with HCC not amenable to curative therapy were randomized in 1:1:1 ratio to receive TACE+durvalumab+bevacizumab, TACE+durvalumab, or TACE+placebo
- Median progression free survival was improved by 6.8 months in the D+B+TACE vs. placebo+TACE arm (15 months vs 8.2 months)

## Y-90 Induces Immune Response



Y90 causes an altered adaptive and innate immune response, resulting in increased frequency of activated CD3+ T Cells and CD8+ regulatory T cells



Increased inflammatory (PD-L1+ and HLA-DR+) monocyte populations





Effect peaks at 1 month after treatment and decreased significant at 3 and 6 months



Using immunotherapy within 1 month post-Y90 could have a synergistic immune response


HCC patients (n=49) with preserved hepatic function (Child Pugh  $\leq$  B7, MELD score  $\leq$  10) and no indication to liver transplantation, undergoing Y<sup>90</sup>TARE (as first-line loco-regional treatment) and longitudinal blood immune monitoring.



Rivoltini L, Bhoori S, Camisaschi C, et al. Y<sup>90</sup>-radioembolisation in hepatocellular carcinoma induces immune responses calling for early treatment with multiple checkpoint blockers *Gut* 2023;72:406-407. CC BY-NC 4.0

# Liver Transplant for HCC

- Historically, candidacy for transplanting patients with HCC has been based off tumor size alone
- Metroticket 2.0 showed that biomarkers and size both play an important role in predicting post-transplant HCC recurrence
- Novel tumor biomarkers such as DCP and AFP-L3 have now been shown to be significantly associated with high-risk explant features



Mazzaferro, Gastroenterology, 2018

# Post Transplant Recurrence




Morphology  
and Biology



Size

# Pushing the Boundaries



Abdelrahim M, Esmail A. Transplant Oncology: An Emerging Discipline of Cancer Treatment. Cancers (Basel). 2023;15(11):2923-2938.

# Autoimmune Hepatitis



PREVALENCE OF 31.2 PER 100,000 PERSONS



FEMALE PREDOMINANCE (4:1)



ONSET PEAKS AT AGES 1-30 AND 40-60




OCCURS FREQUENTLY WITH CONCOMITANT AUTOIMMUNE DISEASES

# Clinical Presentation

- Symptomatic
  - Most patients with AIH present after the development of chronic nonspecific symptoms (fatigue, malaise, arthralgias, or amenorrhea)
  - Easy fatigability is the main complaint in 85% of patients
- Asymptomatic
  - Asymptomatic in 25%-34% of patients
  - Asymptomatic patients infrequently achieve spontaneous laboratory improvement (12%)
  - The absence of symptoms should not discourage treatment
  - Histology similar to symptomatic patients


# Management

## First-Line Treatment of AIH



Mack, AASLD practice guidance, August 2020

# Treatment Options



# Patient Perspective of Treatment


- How satisfied are you with your current AIH treatment?
  - 38% somewhat satisfied
  - 14% dissatisfied
  - 5% very dissatisfied
- What would you most like to change about your treatment of AIH?
  - 56%: Treatment should have less side effects

Medication	Number ever taken	Number discontinued (%)	Most cited reasons for stopping (n, %)
Azathioprine (AZA)	199	57 (29%)	Side effects (30/44, 68%), Toxic metabolism 5/44, 11%)
Mercaptopurine	23	11 (48%)	Toxic metabolism 5/10, 50%)
Mycophenolate mofetil (MMF)	42	7 (17%)	Side effects (4/5, 80%)

Lloyd 2023

# MMF: Mechanism of Action

- MMF is labeled for use in preventing rejection after solid organ transplantation
- MMF is a prodrug of mycophenolic acid
- Inhibits the activity of the type II isoform of inosine-5'-monophosphate dehydrogenase
- Type II isoform is present in immune cells
- Selectively suppresses both T- and B-cell lymphocyte proliferation
- MMF also inhibits monocytes



Research Article  
DLI, Autoimmune, Cholestatic and Genetic Diseases


JOURNAL  
OF HEPATOLOGY

## An open-label randomised-controlled trial of azathioprine vs. mycophenolate mofetil for the induction of remission in treatment-naïve autoimmune hepatitis

Romée J.A.L.M. Snijders<sup>1,2,4,1</sup>, Anna E.C. Stoeltinga<sup>2,1</sup>, Tom J.G. Gevers<sup>2,4,2,4</sup>, Simon Pape<sup>1,2,4</sup>, Maaike Biewenga<sup>2</sup>, Maarten E. Tushuizen<sup>2</sup>, Robert C. Verdonk<sup>2</sup>, Hendrik J.M. de Jonghe<sup>2</sup>, Jan Maarten Wrolijk<sup>2</sup>, Sjoerd F. Bakker<sup>2</sup>, Thomas Vanvoolingheem<sup>2</sup>, Yrjo S. de Boer<sup>1,2,4</sup>, Martine A.M.C. Baven Pronk<sup>1</sup>, Ulrich Beuers<sup>1,2,4</sup>, Adriaan J. van der Meer<sup>1,2</sup>, Nicole M.F. van Gerven<sup>1,2</sup>, Marlijn G.M. Sijtsma<sup>1,2</sup>, Brechje C. van Eijk<sup>1,2</sup>, Manon C. van IJzendoorn<sup>1,2</sup>, Margot van Herwaarden<sup>1,2</sup>, Floris F. van den Brand<sup>1,2</sup>, Kerem Sebti Korkmaz<sup>2</sup>, Aad P. van den Berg<sup>1,2,4</sup>, Maureen M.J. Guichelaar<sup>2</sup>, Amar D. Levens<sup>2</sup>, Bart van Hoek<sup>2</sup>, Joost P.H. Drenth<sup>1,2,4</sup>, on behalf of the Dutch Autoimmune Hepatitis Working Group

Journal of Hepatology 2024, vol. 80 | 576–585

Check for updates




- Overall, 37 (94.9%) patients in the MMF group and 23 patients (74.2%) in the azathioprine group completed treatment
- At week 24, the proportion of patients with biochemical remission was 56.4% in the MMF group (22 of 39 patients) vs. 29.0% in the azathioprine group (9 of 31 patients) (95% CI, 4.0 to 46.7; p = 0.022)
- This difference was also observed in the analysis utilizing only the data available at the 24-week timepoint (p = 0.031).
- Two patients (5.1%) in the MMF group and eight patients (25.8%) in the azathioprine group discontinued treatment owing to AEs/SAEs

Romée J.A.L.M. Snijders, Anna E.C. Stoeltinga, Tom J.G. Gevers, Simon Pape, et al. Published by Elsevier B.V. on behalf of European Association for the Study of the Liver. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). J. Hepatol. 2024, 80, 576–585 (CC BY 4.0)

# MMF vs Azathioprine

## MMF

- Side effect profile: MMF better tolerated than AZA
- Rapidity of action: MMF >> AZA
- Cirrhosis: MMF fewer side effects than AZA
- Works better (?)



## AZA

- Dosing: Aza once daily > MMF BID
- Cost: Aza cheaper than MMF
- Reproduction: AZA safe in pregnancy, MMF teratogenic

# Take Home Points

- Early diagnosis of steatotic liver disease is critical to altering the natural course of the disease process
- New pharmacologic therapy for MASLD/MASH!
- Combination therapy for HCC is a wave of the future
- Expanding liver transplant offers more patients curative treatment
- Consider mycophenolate mofetil for treatment of autoimmune hepatitis

# References

- Byrne CD, Targher G, Tilg H, Thyroid hormone receptor-beta agonists: new MASLD therapies on the horizon *Gut* 2024;73:573-581.
- Cheng AL et al. Updated efficacy and safety data from IMbrave150: Atezolizumab plus bevacizumab vs. sorafenib for unresectable hepatocellular carcinoma. *J Hepatol*. 2022 Apr;76(4):862-873. doi: 10.1016/j.jhep.2021.11.030. Epub 2021 Dec 11. PMID: 34902530.
- Estes, C., Razavi, H., Loomba, R., Younossi, Z. and Sanyal, A.J. (2018), Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology*, 67: 123-133. <https://doi.org/10.1002/hep.29466>
- Finn RS et al. IMbrave150 Investigators. Atezolizumab plus Bevacizumab in Unresectable Hepatocellular Carcinoma. *N Engl J Med*. 2020 May 14;382(20):1894-1905. doi: 10.1056/NEJMoa1915745. PMID: 32402160.
- Guo, X.; Yin, X.; Liu, Z.; Wang, J. Non-Alcoholic Fatty Liver Disease (NAFLD) Pathogenesis and Natural Products for Prevention and Treatment. *Int. J. Mol. Sci.* 2022, 23, 15489. <https://doi.org/10.3390/ijms232415489>
- Harrison SA, Bedossa P, Guy CD, Schattenberg JM, Loomba R, Taub R, Labriola D, Moussa SE, Neff GW, Rinella ME, Anstee QM, Abdelmalek MF, Younossi Z, Baum SJ, Francque S, Charlton MR, Newsome PN, Lanthier N, Schiefke I, Mangia A, Pericas JM, Patil R, Sanyal AJ, Noureddin M, Bansal MB, Alkhouri N, Castera L, Rudraraju M, Ratziu V; MAESTRO-NASH Investigators. A Phase 3, Randomized, Controlled Trial of Resmetrom in NASH with Liver Fibrosis. *N Engl J Med*. 2024 Feb 8;390(6):497-509. doi: 10.1056/NEJMoa2309000. PMID: 38324483.
- *It took 80 horses to film that "Game of Thrones" battle scene.* Horse Network. (2017, September 6). <https://horsenetwork.com/2016/06/it-took-80-horses-to-film-that-game-of-thrones-battle-scene/>
- Lee YB, Nam JY, Cho EJ, Lee JH, Yu SJ, Kim HC, Paeng JC, Yoon JH, Kim YJ. A Phase III Trial of Yttrium-90 Radioembolization in Combination with Durvalumab for Locally Advanced Unresectable Hepatocellular Carcinoma. *Clin Cancer Res*. 2023 Sep 15;29(18):3650-3658. doi: 10.1158/1078-0432.CCR-23-0581. PMID: 37459133.
- Lloyd C, Leighton J, Wong LL, Goulding A, Brownlee A, Gray P, Culver E, Halliday N, Thorburn D, Heneghan MA, Jones DEJ, Eley C, Dyson JK. Patient Priorities in Autoimmune Hepatitis: The Need for Better Treatments, More Education and Challenging Stigma. *Dig Dis Sci*. 2023 Jan;68(1):87-97. doi: 10.1007/s10620-022-07525-5. Epub 2022 May 17. PMID: 35579795; PMCID: PMC9112273.
- Mack, Cara L.<sup>1</sup>; Adams, David<sup>2</sup>; Assis, David N.<sup>3</sup>; Keekar, Nanda<sup>4</sup>; Manns, Michael P.<sup>5</sup>; Mayo, Marilyn J.<sup>6</sup>; Vierling, John M.<sup>7</sup>; Alsawas, Mouaz<sup>8</sup>; Murad, Mohammad H.<sup>9</sup>; Czaja, Albert J.<sup>10</sup>. Diagnosis and Management of Autoimmune Hepatitis in Adults and Children: 2019 Practice Guidance and Guidelines From the American Association for the Study of Liver Diseases. *Hepatology* 72(2):p 671-722, August 2020. | DOI: 10.1002/hep.31065
- Mazzaferro V, Sposito C, Zhou J, Pinna AD, De Carlis L, Fan J, Cescon M, Di Sandro S, Yi-Feng H, Lauterio A, Bongini M, Cucchetti A. Metroticket 2.0 Model for Analysis of Competing Risks of Death After Liver Transplantation for Hepatocellular Carcinoma. *Gastroenterology*. 2018 Jan;154(1):128-139. doi: 10.1053/j.gastro.2017.09.025. Epub 2017 Oct 5. PMID: 28989060.

# References Cont'd

- Newsome PN, Buchholtz K, Cusi K, Linder M, Okanoue T, Ratziu V, Sanyal AJ, Sejjing AS, Harrison SA; NN9931-4296 Investigators. A Placebo-Controlled Trial of Subcutaneous Semaglutide in Nonalcoholic Steatohepatitis. *N Engl J Med*. 2021 Mar 25;384(12):1113-1124. doi: 10.1056/NEJMoa2028395. Epub 2020 Nov 13. PMID: 33185364.
- Rinella, Mary E.<sup>1</sup>; Neuschwander-Tetri, Brent A.<sup>2</sup>; Siddiqui, Mohammad Shadab<sup>3</sup>; Abdelmalek, Manal F.<sup>4</sup>; Caldwell, Stephen<sup>5</sup>; Barb, Diana<sup>6</sup>; Kleiner, David E.<sup>7</sup>; Loomba, Rohit<sup>8</sup>. AASLD Practice Guidance on the clinical assessment and management of nonalcoholic fatty liver disease. *Hepatology* 77(5):p 1797-1835, May 2023. | DOI: 10.1097/HEP.000000000000323
- Rivoltini L, Bhoori S, Camisaschi C, Bergamaschi L, Lalli L, Frati P, Citterio D, Castelli C, Mazzaferro V. Y<sup>90</sup>-radioembolisation in hepatocellular carcinoma induces immune responses calling for early treatment with multiple checkpoint blockers. *Gut*. 2023 Feb;72(2):406-407. doi: 10.1136/gutjnl-2021-326869. Epub 2022 May 4. PMID: 35508369; PMCID: PMC9872224.
- Salem R, Gordon AC, Mouli S, Hickey R, Kallini J, Gabr A, Mulcahy MF, Baker T, Abecassis M, Miller FH, Yaghai V, Sato K, Desai K, Thornburg B, Benson AB, Rademaker A, Ganger D, Kulik L, Lewandowski RJ. Y90 Radioembolization Significantly Prolongs Time to Progression Compared With Chemoembolization in Patients With Hepatocellular Carcinoma. *Gastroenterology*. 2016 Dec;151(6):1155-1163.e2. doi: 10.1053/j.gastro.2016.08.029. Epub 2016 Aug 27. PMID: 27575820; PMCID: PMC5124387.
- Younossi, Z., Tacke, F., Arrese, M., Chander Sharma, B., Mostafa, I., Bugianesi, E., Wai-Sun Wong, V., Yilmaz, Y., George, J., Fan, J. and Vos, M.B. (2019). Global Perspectives on Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Hepatology*, 69: 2672-2682. <https://doi.org/10.1002/hep.30251>