



Gastroesophageal Reflux

Subhankar Chakraborty, M.D., Ph.D.

*Assistant Professor, Division of Gastroenterology, Hepatology & Nutrition
Department of Internal Medicine
The Ohio State University Wexner Medical Center*

MedNet21
Center for Continuing Medical Education

 **THE OHIO STATE UNIVERSITY**
WEXNER MEDICAL CENTER

Objectives

1. Describe the epidemiology, pathophysiology, and risk factors for GERD
2. Recognize the typical and atypical clinical presentations of GERD
3. Apply a stepwise diagnostic approach to GERD
4. Discuss how to manage GERD
5. Explain the risks and benefits of long-term PPI therapy
6. Identify complications of chronic GERD

Case-1

A 45-year-old woman presents with a 6-month history of **burning retrosternal discomfort** occurring after meals and at night. She reports occasional regurgitation but denies dysphagia, odynophagia, weight loss, anorexia, vomiting, or gastrointestinal bleeding. She has tried over-the-counter antacids with minimal relief. Physical examination is unremarkable. What is the **most appropriate next step in management?**

- A) Upper endoscopy
- B) Ambulatory pH monitoring
- C) Empirical trial of a proton-pump inhibitor (PPI) for 2 months
- D) Barium swallow
- E) Esophageal manometry

Case-2

A 54-year-old man presents with a 6-month history of **heartburn and regurgitation**. He reports new-onset dysphagia and unintentional weight loss of 10 pounds over the past two months. He has no prior history of gastrointestinal disease and has not tried acid suppression therapy. Physical examination reveals no acute findings.

Which of the following is the **most appropriate next step in management?**

- A) Empirical trial of a proton-pump inhibitor
- B) Ambulatory pH monitoring
- C) Esophageal manometry
- D) Esophagogastroduodenoscopy (EGD)
- E) Barium swallow

Case-3

A 60-year-old White man with a BMI of 32 kg/m² and a 12-year history of chronic **heartburn and regurgitation** presents for routine follow-up. He is a current smoker and has a family history of esophageal adenocarcinoma. He denies dysphagia, weight loss, or gastrointestinal bleeding. Which of the following is the most appropriate next step?


- A) Empirical trial of a proton-pump inhibitor
- B) Ambulatory pH monitoring
- C) Upper endoscopy to screen for Barrett's esophagus
- D) Esophageal manometry
- E) Barium swallow

What is Gastroesophageal Reflux Disease (GERD)

GERD is a chronic disease characterized by


- **Symptoms** from regurgitation of gastric contents into the esophagus
- Supported by **objective evidence** of acid regurgitation

Understanding GERD: Prevalence and Economic Impact




Created with Google Gemini

Typical and Atypical Symptoms of GERD




Created with Google Gemini

What causes GERD



Created with Google Gemini


Dietary and Behavioral Factors Causing GERD



Created with Google Gemini

Uncommon and Rare Causes of GERD

UNCOMMON CAUSES OF GERD: BEYOND THE TYPICAL FACTORS




Created with Google Gemini

Diagnosing GERD


- Endoscopy:
 - Inflammation or esophagitis
 - Barrett's esophagus
 - Peptic stricture
- Esophageal pH monitoring test
 - High acid levels in distal esophagus

The Endoscopic Spectrum of GERD



Created with Google Gemini

Diagnosing GERD: Objective Tests for Acid Reflux



Created with Google Gemini

When to consider objective testing for GERD

- Symptoms uncontrolled by empiric treatment
- Atypical symptoms
- When considering surgery to treat GERD
- Alarm symptoms
 - Dysphagia
 - Unintentional weight loss
 - Symptoms of upper GI bleeding

Management of GERD

Goals of therapy

- Control symptoms
- Prevent or treat complications
 - **Peptic stricture** → Dysphagia
 - **Barrett's esophagus** → Esophageal adenocarcinoma
 - **Esophageal ulcer** → Odynophagia and GI bleeding






Controlling Symptoms of GERD

- Lifestyle measures
- Medications
- Surgery

Lifestyle Measures to Control GERD

LIFESTYLE MEASURES TO CONTROL GERD SYMPTOMS

MEASURES WITH STRONGEST EVIDENCE **DATA-BACKED**

<p>Weight Loss</p>  <ul style="list-style-type: none"> • DO: Reduce BMI. Dose-dependent symptom improvement. • 40% reduction in symptoms with ≥ 3.5 BMI reduction (in women). • DO: Avoid weight gain. • Interventions decreased prevalence 37% \rightarrow 15%. 	<p>Head-of-Bed Elevation</p>  <p>RCT data supports (multiple studies).</p> <p>6-8 inches (15-20 cm)</p> <ul style="list-style-type: none"> • DO: Elevate bed head end. Reduces nocturnal reflux episodes. • DO: Use a wedge pillow. RCT data supports (multiple studies). 	<p>Avoid Late Meals</p>  <p>no food over the last 3 hours before midnight.</p> <ul style="list-style-type: none"> • DON'T: Eat large meals especially at bedtime. • DON'T: Eat within 3 hrs of bedtime. • DO: Avoid short dinner-to-bed intervals. 	<p>Smoking Cessation</p>  <p>44% improvement</p> <ul style="list-style-type: none"> • DO: Stop smoking. • 44% symptom improvement after 1 year. • Validated data supports.
OTHER IMPORTANT LIFESTYLE CHANGES			
<p>Sleep Positioning</p>  <p>LEFT SIDE preferential (reflux reduced) RIGHT SIDE dependent position (increased reflux)</p> <p>DO: Sleep in left lateral position. Positional pillows help. DON'T: Sleep on right side.</p>	<p>Dietary Modifications</p> <p>MEAL SIZE 4 small meals vs 3 large meals</p> <p>AVOID TRIGGERS Spicy/Fatty Foods, Onion, Citrus, Chocolate</p> <p>LIMIT BEVERAGES Coffee, Tea, Carbonated Drinks</p> <p>Water substitution</p> <p><small>Nurses Health Study data</small></p> <p><small>ASGE: Alcohol cessation data is limited for GERD specifically, though not recommended.</small></p>	<p>Behavioral Measures</p> <p>CLOTHING vs DON'T</p> <p>BENDING vs DON'T</p> <p>CHEW GUM</p> <p>MEDICATIONS Benzos, CCBs, Antidepressants etc.</p> <p>STRESS REDUCTION DO: Relaxation, Breathing, CBT. Leverages Gut-Brain Axis.</p>	

Created with Google Gemini

Proton Pump Inhibitors

- Highly effective in healing esophagitis: 84% heal within 3 months compared to 52% for H2-blockers (e.g. Famotidine)
- Duration
 - **Short-term PPI (4-8 weeks):** Uncomplicated GERD
 - **Long-term PPI:**
 - Barrett's esophagus
 - Severe esophagitis
 - Complicated GERD
 - **On-demand PPI:** episodic heartburn and GERD symptoms without inflammation

Proton Pump Inhibitors

- **Administration:** 30-45 mins before 1st meal of the day or before the evening meal if taking BID
- **Duration:** At least 4-8 weeks for patients without alarm symptoms
- **Discontinuation:** If symptoms improve, decrease the dose and ultimately discontinue
- Any PPI will do- no one is superior to another










Long-Term Risks of PPI Use

DEFINITIVE RISK: SMALL INCREASE IN ENTERIC INFECTIONS

Non-*C. difficile* Enteric Infections

- **DO:** Be aware of a small definitive risk.
- **Odds Ratio (OR):** 1.33 (1.01–1.75)
- Absolute risk is **VERY SMALL** (0.03–0.2% per patient/year)
- Includes bacterial gastroenteritis, bacterial overgrowth

OBSERVATIONAL ASSOCIATIONS: RISKS ARE NOT SIGNIFICANTLY INCREASED

 <p>Clostridium difficile Infection OR: 2.26 (0.70–7.34) Non-significant association</p>	 <p>Chronic Kidney Disease OR: 1.17 (0.94–1.45) Non-significant association</p>	 <p>Dementia OR: 1.20 (0.81–1.78) Non-significant association</p>
 <p>Bone Fracture OR: 1.02 (0.87–1.19) Non-significant association</p>	 <p>Myocardial Infarction Not Significantly Increased</p>	 <p>SBP in Cirrhotics Reported in observational studies; not significant</p>
 <p>Micronutrient Deficiencies Not Significantly Increased</p>	 <p>Reported in observational studies; not significant</p>	 <p>GI Malignancies Reported in observational studies; not significant</p>

Current evidence does not support significant increases in most widely studied associations.
Close patient monitoring remains important.

Created with Google Gemini

Short Term Side Effects of PPIs

Most Common Adverse Reactions

Mild and resolve with discontinuation, may be PPI preparation specific

- Headache
- Abdominal pain
- Nausea or vomiting
- Diarrhea or Constipation
- Flatulence

Rebound Acid Hypersecretion

- Upon PPI discontinuation
- Due to PPI-induced hyperplasia of gastric enterochromaffin-like cells that release gastrin
- Can be mitigated by gradual tapering of PPI dose

Other Side Effects of PPIs

- Acute interstitial Nephritis
- Hypomagnesemia
- Fundic gland polyps

Created with Google Gemini

Other Side Effects of PPIs

- **Acute interstitial Nephritis**
 - Can occur at any time and at any dose
 - Suspect if unexplained rise in serum creatinine, particularly if nonoliguric
 - Stop PPI immediately
- Hypomagnesemia
- Fundic gland polyps

Created with Google Gemini

Other Side Effects of PPIs

- Acute interstitial Nephritis
- **Hypomagnesemia**
 - Risk factors:
 - high dose PPI
 - concomitant diuretics
 - low dietary magnesium
 - Females
 - Diabetics
 - CKD
- Fundic gland polyps

Created with Google Gemini

Other Side Effects of PPIs

- Acute interstitial Nephritis
- Hypomagnesemia
- **Fundic gland polyps**
 - More likely with longer duration of PPI use
 - Very low risk of dysplasia (1%) compared to FGPs associated with Familial Adenomatous Polyposis (25-46% dysplasia)
 - Routine endoscopic surveillance not recommended
 - Solitary polyps and large FGPs should be resected

Created with Google Gemini

Potassium Competitive Acid Blockers

- **Vonoprozán-** available in USA
- **Administration:** Before or After meals
- **Efficacy in healing inflammation:** More than 80%
- **Duration:** At least 4-8 weeks for patients without alarm symptoms
- **Discontinuation:** If symptoms improve, can discontinue
- **Risks:** Similar to PPIs

Created with Google Gemini

Histamine-2 Receptor Antagonists

- **Famotidine-** available in the USA
- **Administration:** before or after meals
- **Efficacy in healing inflammation:** Less effective
- **Duration:** As needed for patients without alarm symptoms
- **Risks:** Tachyphylaxis

Created with Google Gemini

Antacids, Alginates and Mucosal Protectants

- Provide short-term symptomatic relief
- Useful for nocturnal or post-meal symptoms
- Used as adjuncts to PPIs in those with erosive esophagitis
- **Antacids:** Calcium carbonate
- **Alginates:** Reflux Gourmet™, Gaviscon Advance™
- **Mucosal protectant:** Sucralfate (tablet or Suspension)

Created with Google Gemini

Baclofen

- **Mechanism:** Reduces the frequency of transient lower esophageal sphincter relaxation
- **Application:** Adjunct in regurgitation predominant GERD or Belching
- **Side effects:** Dizziness, fatigue, sleepiness

Created with Google Gemini

Surgery For Reflux

- When:
 - Refractory symptoms despite optimum medical therapy
 - Patient's desire to avoid long term medications
 - Large hiatal hernias
- What
 - Fundoplication
 - Transoral Incisionless Fundoplication (TIF)
 - Magnetic sphincter augmentation
 - Roux-en Y gastric bypass







Created with Google Gemini

Surgery For Reflux

- When not to recommend surgery:
 - Patient well controlled on medical therapy
 - To prevent esophageal cancer in those with Barrett's esophagus

Created with Google Gemini

Anti-Reflux Interventions: Key Features and Candidate Selection







<p style="text-align: center;">NISSEN (360° total fundoplication)</p>  <p>Complete posterior wrap (360°) around distal esophagus</p> <p>IDEAL CANDIDATE Normal esophageal motility; severe GERD (LA C/D); large hiatal hernia</p>	<p style="text-align: center;">TOUPET (270° posterior)</p>  <p>Posterior partial wrap (270°) leaving anterior esophagus uncovered</p> <p>IDEAL CANDIDATE Impaired esophageal motility or borderline peristaltic reserve; concern for postoperative dysphagia</p>
<p style="text-align: center;">DOR</p>  <p>Anterior partial wrap (180°) over esophagus</p> <p>IDEAL CANDIDATE Primarily after Heller myotomy for achalasia; rarely used as primary GERD procedure</p>	<p style="text-align: center;">MSA (LINX device)</p>  <p>Ring of magnetic beads augmenting LES</p> <p>IDEAL CANDIDATE Non-obese; regurgitation-predominant; small hiatal hernia (≤3 cm); no prior gastric surgery</p>
<p style="text-align: center;">TIF 2.0 (endoscopic)</p>  <p>Transoral plication creating 270° flap valve at EGJ</p> <p>IDEAL CANDIDATE No/small hiatal hernia (≤2 cm); BMI < 35; troublesome regurgitation; patients declining surgery</p>	<p style="text-align: center;">RYGB</p>  <p>Small gastric pouch and long Roux limb prevents bile reflux</p> <p>IDEAL CANDIDATE Obese patients (BMI ≥35) with GERD; failed fundoplication in obese; concurrent metabolic disease</p>

These features help determine the most suitable intervention for each patient, based on comprehensive selection criteria. Close patient monitoring remains important.

Created with Google Gemini

Anti-Reflux Interventions: Key Advantages and Disadvantages

SURGICAL & ENDOSCOPIC OPTIONS FOR GERD: PROS AND CONS (SIMPLE SUMMARY)


NISSEN FUNDOPPLICATION (360°, total wrap)	TOUPET FUNDOPPLICATION (270°, posterior partial wrap)	DOR FUNDOPPLICATION (180°, anterior partial wrap)	LINX (Magnetic sphincter augmentation)	TIF 2.0 (Endoscopic fundoplication)	ROUX-EN-Y GASTRIC BYPASS (RYGB)
 <ul style="list-style-type: none"> ✓ Best reflux control (gold standard) ✓ Strong LES pressure increase ✓ Excellent long-term durability (>20 years) ✗ Highest risk of dysphagia (~40% early) ✗ Gas-bloat common (~30%) ✗ Inability to belch or vomit 	 <ul style="list-style-type: none"> ✓ Reflux control similar to Nissen ✓ Much less dysphagia (~10% early) ✓ Preserves esophageal motility ✓ Little to no gas-bloat ✗ Possibly higher long-term GERD recurrence (debated) 	 <ul style="list-style-type: none"> ✓ Less dysphagia than Nissen ✓ Effective reflux control after Heller myotomy ✗ Weaker reflux barrier than Nissen or Toupet ✗ Limited data as primary GERD surgery 	 <ul style="list-style-type: none"> ✓ Minimally invasive ✓ Shorter surgery time ✓ Preserves ability to belch and vomit (~96%) ✓ Reversible ✗ MRI restrictions (>1.5T) ✗ Small erosion risk (~0.3%) ✗ Device removal risk (~4%) ✗ Dysphagia can be higher than fundoplication 	 <ul style="list-style-type: none"> ✓ Incisionless, outpatient procedure ✓ High short-term symptom control ✓ Many patients reduce or stop PPIs ✗ Not for severe esophagitis or large hernias ✗ Long-term durability uncertain ✗ Most patients resume low-dose PPIs 	 <ul style="list-style-type: none"> ✓ Treats both obesity and GERD ✓ Major reduction in reflux symptoms ✓ High PPI discontinuation rates ✗ Major anatomical change ✗ Nutritional deficiencies ✗ Technically complex ✗ Reflux can recur; new GERD possible

✓ Key Advantages ✗ Key Disadvantages

👤 Optimal anti-reflux intervention depends on comprehensive patient selection (confirmed GERD, motility assessment, BMI, hernia size).

Created with Google Gemini


Management of GERD



Created with Google Gemini

Management of GERD

MANAGEMENT OF GERD: STEPS 3 & 4







Created with Google Gemini

Management of GERD

MANAGEMENT OF GERD: STEPS 5 & 6

STEP 5 – PHENOTYPE-DRIVEN ADJUNCTIVE THERAPY


When symptoms persist despite optimized PPIs, adjunctive therapies should match the dominant symptom pattern:


	Post-prandial or breakthrough symptoms → Alginate antacids	
	Nocturnal symptoms → Bedtime H2-receptor antagonists	
	Regurgitation- or belch-predominant symptoms → Baclofen	
	Coexistent gastroparesis → Prokinetics	

Internist takeaway: Not all persistent symptoms mean “more acid.” Target the physiology.

STEP 6 – FUNCTIONAL ESOPHAGEAL DISORDERS

Patients with:


✓ Normal endoscopy 


✓ Physiologic acid exposure 

Symptoms triggered by reflux events or unrelated to reflux.

Often have: Functional heartburn, or Reflux hypersensitivity.

Recommended approach:

 Neuromodulators (e.g., low-dose antidepressants)

 Behavioral therapies (CBT, hypnotherapy, diaphragmatic breathing)

 Gradual PPI de-escalation, as tolerated

Key concept: These patients do not benefit from escalating acid suppression.

Created with Google Gemini

GERD Management in Special Populations



1. PATIENTS WITH EXTRAESOPHAGEAL SYMPTOMS (Chronic cough, laryngitis, asthma)



WHAT TO DO


 Do **NOT** start with empiric PPI therapy

 Order objective reflux testing **OFF** acid-suppressive therapy first

WHY

 PPI response rates are low 

 Most have non-reflux causes 

 Avoids unnecessary long-term PPI use and diagnostic delay

Practical takeaway: If the main complaint is cough or throat symptoms without classic heartburn, test first—do not trial PPIs.


2. OBESE PATIENTS (BMI ≥35)

WHAT TO DO

PREFERRED PROCEDURE: RYGB




WHY

 RYGB improves both obesity and GERD

AVOID SLEEVE



 Sleeve gastrectomy often worsens reflux

Practical takeaway: In patients with severe obesity and GERD, bariatric surgery choice matters—RYGB treats reflux; sleeve may aggravate it.

3. SEVERE EROSIVE ESOPHAGITIS (LA Grade C or D)


WHAT TO DO

 Indefinite maintenance PPI or anti-reflux surgery

WHAT TO DO

 8–12 weeks  Repeat endoscopy in 8–12 weeks after starting treatment

WHY

 High relapse risk if PPIs are stopped

WHY

 Repeat EGD ensures healing and rules out Barrett's esophagus

Practical takeaway: These patients need lifelong reflux control and documented mucosal healing.

Created with Google Gemini

When Should You Refer GERD Patients to a Gastroenterologist?

URGENT REFERRAL: ALARM SYMPTOMS	NON-RESPONDERS & UNCERTAINTY	SCREENING & CONFIRMATION	SPECIAL SITUATIONS & COMPLICATIONS
<ul style="list-style-type: none"> DYSPHAGIA ODYNOPHAGIA WEIGHT LOSS GI BLEEDING IRON-DEFICIENCY ANEMIA PERSISTENT VOMITING <p>NEEDS PROMPT UPPER ENDOSCOPY</p>	<p>REFRACTORY SYMPTOMS</p> <ul style="list-style-type: none"> Persistent symptoms despite optimized PPI Correct timing & adherence BID PPI for ~8 weeks Recurrence after stopping PPI <p>DIAGNOSTIC UNCERTAINTY</p> <ul style="list-style-type: none"> Need to confirm or exclude GERD Requires specialized testing: <ul style="list-style-type: none"> Upper Endoscopy Ambulatory pH/pH-Impedance monitoring 	<p>BARRETT'S ESOPHAGUS SCREENING</p> <ul style="list-style-type: none"> Patients meeting guideline risk thresholds (e.g., chronic GERD with multiple additional risk factors) <p>OBJECTIVE CONFIRMATION</p> <ul style="list-style-type: none"> Before long-term PPI use For patients started on chronic PPIs without proven GERD Prove GERD with Endoscopy & Reflux Testing first 	<p>ISOLATED EXTRAESOPHAGEAL SYMPTOMS</p> <ul style="list-style-type: none"> Chronic cough, laryngitis, asthma with suspected reflux Requires UPFRONT OBJECTIVE REFLUX TESTING (not empiric PPI trial) <p>CONSIDERING ANTI-REFLUX PROCEDURES</p> <ul style="list-style-type: none"> Evaluation for: <ul style="list-style-type: none"> Fundoplication LINX Endoscopic therapy Requires specialized testing (Endoscopy, Manometry, Reflux Monitoring) <p>COMPLICATED GERD</p> <ul style="list-style-type: none"> Needs specialized management LA Grade C/D esophagitis Peptic strictures Barrett's esophagus Requires healing confirmation, surveillance, endoscopic therapy

Created with Google Gemini

Screening for Barrett's Esophagus

WHO SHOULD BE SCREENED FOR BARRETT'S ESOPHAGUS?

CORE PRINCIPLE: SCREEN WHEN RISK ACCUMULATES, NOT BASED ON REFLUX SYMPTOMS ALONE.

Screening is best reserved for patients with multiple established risk factors, where likelihood of BE and EAC is meaningfully higher.

CONSIDER SCREENING

ENDOSCOPY IN PATIENTS WITH CHRONIC GERD PLUS ADDITIONAL RISK FACTORS, ESPECIALLY WHEN SEVERAL ARE PRESENT.

Screening is not recommended for everyone with GERD.

CONSISTENTLY RECOGNIZED RISK FACTORS

Male sex	Age >50 years	White / non-Hispanic white race
Chronic GERD ≥5 YEARS OF SYMPTOMS OR WEEKLY HEARTBURN OR REGURGITATION	Obesity, especially central (abdominal) obesity	Current or prior tobacco
	Obesity, especially central (abdominal) obesity	First-degree family history First-degree family history of Barrett's esophagus or esophageal adenocarcinoma

HOW RISK FACTORS ADD UP: VISUALIZING ACCUMULATING RISK

CLINICAL IMPLICATION: RISK FACTORS ARE ADDITIVE, AND SCREENING YIELD RISES SHARPLY AS THEY ACCUMULATE.

general medical authority

Created with Google Gemini

Case-1

A 45-year-old woman presents with a 6-month history of **burning retrosternal discomfort** occurring after meals and at night. She reports occasional regurgitation but denies dysphagia, odynophagia, weight loss, anorexia, vomiting, or gastrointestinal bleeding. She has tried over-the-counter antacids with minimal relief. Physical examination is unremarkable. What is the most appropriate next step in management?

- A) Upper endoscopy
- B) Ambulatory pH monitoring
- C) **Empirical trial of a proton-pump inhibitor (PPI) for 2 months**
- D) Barium swallow
- E) Esophageal manometry

Answer: **Empirical trial of a proton-pump inhibitor (PPI) for 2 months.**

Rationale: In the absence of alarm symptoms (such as dysphagia, odynophagia, weight loss, anorexia, vomiting, or gastrointestinal bleeding), an empirical course of PPI therapy is a reasonable initial diagnostic and therapeutic approach for patients with typical GERD symptoms. This strategy is supported by evidence showing that a short course of high-dose PPI has a pooled sensitivity of 79% for GERD diagnosis in patients with typical symptoms and no alarm features. Upper endoscopy is reserved for patients with alarm symptoms or those at risk for Barrett's esophagus.

Fass R., New England Journal of Medicine [Volume 387 • Number 13 • September 29, 2022](#); Pages: 1207-1216

Case-2

A 54-year-old man presents with a 6-month history of heartburn and regurgitation. He reports new-onset dysphagia and unintentional weight loss of 10 pounds over the past two months. He has no prior history of gastrointestinal disease and has not tried acid suppression therapy. Physical examination reveals no acute findings.

Which of the following is the most appropriate next step in management?

- A) Empirical trial of a proton-pump inhibitor
- B) Ambulatory pH monitoring
- C) Esophageal manometry
- D) **Esophagogastroduodenoscopy (EGD)**
- E) Barium swallow

Answer: **Esophagogastroduodenoscopy (EGD).**

Rationale: Upper endoscopy is indicated in patients with GERD symptoms who present with alarm features such as dysphagia or weight loss, as these may signal underlying complications including malignancy or strictures. EGD allows direct visualization and biopsy of the esophageal mucosa to evaluate for erosive esophagitis, Barrett's esophagus, or other pathology.

Fass R., New England Journal of Medicine [Volume 387 • Number 13 • September 29, 2022](#); Pages: 1207-1216

Case-3

A 60-year-old White man with a BMI of 32 and a 12-year history of chronic heartburn and regurgitation presents for routine follow-up. He is a current smoker and has a family history of esophageal adenocarcinoma. He denies dysphagia, weight loss, or gastrointestinal bleeding. Which of the following is the most appropriate next step?

- A) Empirical trial of a proton-pump inhibitor
- B) Ambulatory pH monitoring
- C) **Upper endoscopy to screen for Barrett's esophagus**
- D) Esophageal manometry
- E) Barium swallow

Answer: Upper endoscopy to screen for Barrett's esophagus is indicated in patients with chronic (≥ 5 years) GERD symptoms and three or more of the following risk factors: male sex, age > 50 years, White race, tobacco smoking, obesity, or a family history of Barrett's esophagus or esophageal adenocarcinoma. This approach is justified because these factors are associated with increased risk for Barrett's esophagus and esophageal adenocarcinoma, and targeted screening may allow for earlier detection and management of premalignant changes

Fass R., New England Journal of Medicine [Volume 387 • Number 13 • September 29, 2022](#); Pages: 1207-1216

Summary: Take Home Points

- Optimize PPI timing, dose, and adherence
- Many PPI nonresponders lack true GERD
- Use objective testing before long-term PPI therapy
- Titrate PPIs to the lowest effective dose
- Match adjunctive therapy to symptom phenotype
 - Alginates for postprandial breakthrough
 - Famotidine for nocturnal symptoms
 - Baclofen for regurgitation or belch-predominant symptoms.
- Confirm reflux before surgical referral
- Procedure choice depends on anatomy and BMI

References

- **AGA CPU on GERD Personalized Evaluation and Management.** *Clin Gastroenterol Hepatol.* 2022. Yadlapati R, Gyawali CP, Pandolfino JE.
- **ACG Clinical Guideline: Diagnosis and Management of GERD.** *Am J Gastroenterol.* 2022. Katz PO, Dunbar KB, Schnoll-Sussman FH, et al.
- **Gastroesophageal Reflux Disease.** *Yamada's Textbook of Gastroenterology, 7th ed.* 2022. Roman S, Pandolfino JE, Kahrilas PJ.
- **Gastroesophageal Reflux Disease.** *Evidence-Based Gastroenterology and Hepatology, 4th ed.* 2019. Roman S, Kahrilas PJ.
- **Refractory Heartburn: Reflux Hypersensitivity and Functional Heartburn.** *The Esophagus.* 2021. Shibli F, Fass R.
- **AGA CPU on Screening and Surveillance Innovation in Barrett's Esophagus.** *Clin Gastroenterol Hepatol.* 2022. Muthusamy VR, Wani S, Gyawali CP, et al.
- **ACG Guideline: Diagnosis and Management of Barrett's Esophagus.** *Am J Gastroenterol.* 2022. Shaheen NJ, Falk GW, Iyer PG, et al.
- **ASGE Guideline on Barrett's Esophagus Screening and Surveillance.** *Gastrointest Endosc.* 2019. Qumseya B, Sultan S, Bain P, et al.
- **Barrett Esophagus: Rapid Evidence Review.** *Am Fam Physician.* 2022. Bryce C, Bucaj M, Gazda R.
- **Oesophageal Cancer.** *Lancet.* 2024. Yang H, Wang F, Hallemeier CL, et al.
- **Esophageal and Esophagogastric Junction Cancers.** *NCCN Guidelines.* Updated Jan 21, 2026.