




Autism Spectrum Disorder

Christopher Hanks, MD
*Associate Professor - Clinical
 Department of Internal Medicine
 Division of General Internal Medicine and Geriatrics
 The Ohio State University Wexner Medical Center*


MedNet21
Center for Continuing Medical Education

THE OHIO STATE UNIVERSITY
WEXNER MEDICAL CENTER



Disclosures

- Dr. Hanks receives grant funding from the following sources:
 - Bill and Marci Ingram & The White Castle Foundation
 - Operational funding for The Center for Autism Services and Transition
 - Department of Defense AWD-112301
 - Promoting Residents' Excellence in Patient-centered cARE (PREPARE) for autistic adults



Learning Objectives

- Understand the core features of autism and how they present throughout the life course
- Learn the impact of the medical vs social model of disability on autistic individuals
- Recognize common co-occurring diagnoses
- Increase comfort in recognition and management of physical, mental, and behavioral concerns that may occur in autistic individuals

What is autism?¹

- Biologically based
- Diagnosis based on
 - Difference in social communication and interaction
 - Repetitive or restrictive patterns of behavior, interests, or activities
- Signs and Symptoms are present in early childhood
 - May not become obvious until later in childhood, or adulthood, when demands increase

Etiology of autism?²⁻⁵




- Genetic factors
 - 40-90% heritability rate
 - Most autistic people do not have monogenic findings
 - Reasonable to offer genetic testing to any individual diagnosed as autistic.
- Prenatal/Perinatal exposures?
- No evidence to support vaccines causing autism

Prevalence of autism?

Prevalence of Autism Spectrum Disorder
Data from ADDM Network 2000-2018 ⁶

Most recent CDC estimates:
• 1 in 44 children
• 2.2% of adults



Medical vs Social Model of Disability


Medical Model

- Focused on the individual and
 - Their impairment
 - Remediation of skills so that an individual can progress along a typical developmental pathway
- The source of words like "disorder" and "deficit"
- The search for a cure for autism

Social Model


- Views the disability as a result of:
 - Environmental barriers
 - Societal barriers
- Looks at not just impairment, but also strengths and abilities
- Neurodiversity movement

Language of autism⁷



Screening for autism


- MCHAT-R/F⁸: age 16-30 months
 - Sensitivity 0.83
 - Specificity 0.94
- ASSQ⁹: age 7-16 years
 - Sensitivity 91%
 - Specificity 86%
- AQ¹⁰: adults
 - Sensitivity 79%
 - Specificity 98%



The diagram shows a process for the diagnosis of autism. At the top, four icons represent a stethoscope, test tubes, an hourglass, and a thumbs-up. Below these is the title "Diagnosis of Autism". A box on the left titled "Comprehensive Evaluation" lists "History and physical" and "Diagnostic tools: ADOS, ADI-R, CARS-2, other". A double-headed arrow connects this box to a box on the right that states "No need to wait for diagnosis to refer for intervention".

The Role of Primary Care for Autistic Patients

1. Avoid diagnostic overshadowing
2. Age-appropriate health care
3. Accommodate for individual needs
4. Refer to appropriate resources



Preventative Healthcare and Age-appropriate care

- Age-appropriate screenings
 - Cancer screenings
 - Labs
 - Screening for substance abuse
 - STI screening
- Vaccinations
- Other

Providing appropriate accommodations

Clinical settings with systems focused on accommodations can result in¹²⁻¹⁵:

- Higher likelihood of preventative healthcare services
- Higher levels of satisfaction of care
- Fewer unmet healthcare needs
- Higher continuity of care
- Lower expenditures for mental health admissions
- Fewer inpatient hospitalizations and ED visits

Referring to appropriate supports and resources

- Therapies: ST, OT, PT, Behavioral Health, ABA, other
 - Early intervention, School-based, Private, etc
- Physician specialists
- Social supports:
 - Board of developmental disabilities
 - School-based
 - Vocational supports
- Key ages to think about
 - Under 3 years – early intervention
 - 18 years – supported decision-making
 - 22 years – end of school-based special-education supports
 - 26 years – change from parent-based insurance

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Arlington: American Psychiatric Association; 2013.
2. Bai D et al. Association of Genetic and Environmental Factors with autism in a 5-county cohort. *JAMA Psychiatry*. 2019;76(10):1035.
3. Hallmayer J, Cleveland S, Torres A, Phillips J, Cohen B, Torigoe T, et al. Genetic heritability and shared environmental factors among twin pairs with autism. *Arch Gen Psychiatry*. 2011;68(11):1095–102.
4. Grafodatskaya D, Chung B, Szatmari P, Weksberg R. Autism spectrum disorders and epigenetics. *J Am Acad Child Adolesc Psychiatry*. 2010;49(8):794–809.
5. Taylor SE, Way BM, Welch WT, Hilmert CJ, Lehman BJ, Eisenberger NI. Early family environment, current adversity, the serotonin transporter promoter polymorphism, and depressive symptomatology. *Biol Psychiatry*. 2006;60(7):671–6.
6. Source: <https://www.cdc.gov/ncbddd/autism/data.html> – viewed on 2/15/23 – ADDM – CDC’s Autism and Developmental Disabilities Monitoring Network.
7. Bottema-Beutel K, Kapp SK, Lester JN, Sasson NJ, Hand HN. Avoiding ableist language: suggestions for autism researchers. *Autism in Adulthood*. 2021;3(1):18–29.
8. Truhanova Wiczkowski A, et al. Sensitivity and Specificity of the modified checklist for autism in toddlers (original and revised). *JAMA Pediatr*. Published online Feb 20, 2023.
9. Fossevad et al. Validation of the autism spectrum screening questionnaire in a total population sample. *J Autism Dev Disord* 2009;39(1):126–34
10. Baron-Cohen S, et al. The autism-spectrum quotient: evidence from asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *J Autism Dev Disord* 2001;31(1):15–17
11. Croen L, et al. The health status of adults on the autism spectrum. *Autism* 2015;19(7):814–823
12. Hand B, et al. Effects of a specialized primary care facility on preventive service use among autistic adults: a retrospective claims study. *J Gen Intern Med* 2021;36(6):1085–1088
13. Hand B, et al. Patient and caregiver experiences at a specialized primary care center for autistic adults. *J Comp Eff Res* 2020;9(16): 1131–1140
14. Hand B, et al. Specialized primary care medical home: a positive impact on continuity of care among autistic adults. *Autism* 2021;25(1):258–265
15. Alaim, G, et al. Expenditures and Healthcare Utilization of Patients Receiving Care at a Specialized Primary Care Clinic Designed with and for Autistic Adults. *J Gen Int Med* 2022;37(10):2413–2419.



Behavioral and Psychiatric Manifestations of Autism Spectrum Disorder

Ashley Byrd, MD
*Assistant Professor - Clinical
 OSU Department of Psychiatry & Behavioral Health
 Psychiatrist - Nisonger Center - UCEDD
 The Ohio State University Wexner Medical Center*

MedNet21
Center for Continuing Medical Education



Behavioral Manifestations

Restrictive and Repetitive Behaviors (RRBs)

- **Essential feature of ASD**
 - Repetition
 - Inflexibility
 - Invariance
 - Inappropriateness and lack of obvious function/purpose
 - Restricted/fixated interests

Repetitive motor behaviors – stereotypies, rocking, hand flapping, moving fingers in front of eyes, spinning toys, flipping light switches repeatedly

Ritualistic behaviors – insistence on sameness, resistance to change, repetitive language, limited interests, eating same foods, watching same movie (or even parts) repeatedly, resistance to change in environment

Behavioral Manifestations

Restrictive and Repetitive Behaviors (RRBs)

Beneficial vs Impairing

- Pt with fixated interest in weather – pursuing career in meteorology; delays sleep to track storms

Functional impairment

- school performance
- learning essential life skills
- social relationships
- problem behaviors to discourage change/interruption

Behavioral Manifestations

Challenging (Problem) Behaviors

Estimated 50+% of individuals engage in at least one
 They are hard to address, and can significantly affect daily life

- **Aggression**
 - scratching, biting, kicking
 - property destruction
- **Self-injurious behavior**
 - scratching, hair pulling, headbanging
- **Tantrums**
- **Noncompliance**

Behavioral Manifestations
Challenging (Problem) Behaviors

Operant Conditioning ‘Setting events’

Antecedent (stimulus) → Behavior • internal – biological
 → Consequence(s) • external – environmental

Antecedents can trigger, consequences can reinforce behavior

Common internal

- Allergies
- Anxiety
- Constipation/GERD
- Migraines
- Sleep disorders
- Pain

- Escape a demand
- Denial
- Punishment
- Medical

Behavioral Manifestations
Challenging (Problem) Behaviors

Internal Sensory Perception (Interoception)
 Mediated by anterior insula and ventromedial prefrontal cortex
 Studies suggest impaired function in autism

- Hyperresponsive
- Hyporesponsive

Communication Difficulty
 Limited or inability to express wants and needs → frustration → CB

Challenging behaviors are a primary reason individuals come to psychiatric attention

Psychiatric Manifestations
Co-Occurring Psychiatric Conditions

Anxiety Disorders - 40-80%
 ADHD – 30-50%
 Obsessive Compulsive Disorder - ~17%
 Depression – 10-70%
 Bipolar Disorder – 5-8%
 Psychosis - ~35%
 Catatonia – 12-18%

Psychiatric Manifestations
Assessment of Psychiatric Conditions

Features of ASD can overlap with different psychiatric conditions

- Determine patient’s baseline
- Consider developmental level
- Rule out medical conditions
 - hypothyroidism and OSA for depression
- Consider genetics
 - William’s Syndrome – anxiety
 - 22q11 Deletion Syndrome - psychosis

Anxiety Disorders

Risk factors: social skills, sensory sensitivity, rigidity
 Associated with challenging behaviors

Specific Phobia

- Sensory driven
- Noise, needles, large crowds
- Can present as avoidance behaviors

Anxiety Disorders

Generalized Anxiety

- Atypical features
 - preoccupations
 - schedule/change
- Can present as repetitive questions, reassurance seeking about specific events/worries
- Some have difficulty vocalizing - fidgeting, pacing, hand-wringing

Social Anxiety

- Distinguish lack of interest in socialization from avoidance due to fear of embarrassment/judgment

ADHD

Impairments in attention, hyperactivity, and impulsivity

Overlapping features:

- Inattention
- Executive dysfunction
- Social/communication deficits
- Impulsivity
- Restlessness
- Hyperactivity

DSM-IV vs DSM-5

ADHD + ASD

- Increased psychosocial difficulty
- Lower quality of life

Assessment

- Distractibility related to fixated interest, sensory seeking behavior
- Lack of focus due to disinterest rather than concentration
- Excessive talking/interruption - specific interest?

Obsessive Compulsive Disorder

Obsession - Recurrent and persistent thoughts, urges, or images that are experienced... as intrusive and unwanted, and... cause marked anxiety or distress.

Compulsion - Repetitive behaviors (e.g., hand washing, ordering, checking) or mental acts (e.g., praying, counting, repeating words silently) that the individual feels driven to perform in response to an obsession or according to rules that must be applied rigidly.

Overlapping features - RRBs

Key differences:

- OC - distressing, bothersome, unwanted
- ASD - preferred or comforting; consider stereotypies, ritualistic behaviors

Psychotic Disorders

Hallucinations, delusions, disorganized speech, disorganized behaviors

Overlapping features:

- Language pragmatics - disorganized speech
- Under stress - can appear disorganized
- Idiosyncratic/Scripted language - disorganized/delusional
- Restricted interests - delusional
- Theory of Mind - distrust/paranoid
- Self-dialogue - hallucinations

Assessment:

- Developmental history
- Psychotic symptoms usually develop in adolescence or adulthood
- Features above which have been present for years since childhood more likely d/t ASD
- Key is differentiating baseline and assessing change in function

Catatonia

Complex neuropsychiatric behavioral syndrome characterized by abnormal movement, communication, behaviors and withdrawal

- In neurotypical population associated with mental and medical illness
- In ASD can develop in adolescence/early adulthood

Presentation

- Immobility/Stupor
- Mutism
- Staring
- Posturing
- Grimacing
- Echopraxia/echolalia
- Stereotypies
- Withdrawal
- Verbergeration

Catatonia

Overlapping features:

- Echolalia
- Mutism
- Stereotypic movements

Assessment:

- Developmental history, baseline behaviors and assessing for change
- Loss of skills (self care, toileting, etc) - needing prompts or guidance on things previously mastered
- Reduced speech
- Difficulty starting tasks
- Getting "stuck"
- Repetitive behaviors - if present at baseline, have they increased?

Management of Behavioral and Psychiatric Concerns

Behavioral Interventions

Mainstay of treatment

Challenging Behaviors

- Medical exam – acute or chronic illness, pain, etc.
- Psychiatric exam

Applied Behavioral Analysis (ABA)

Most well-known therapy for children with autism

Features

- Social skills, communication, self-care, learning, etc. (individualized)
- School, home, community
- Several hours per week
- Antecedents and Consequences (Functional Behavioral Assessment)
- Goal – to be independent as much as possible; reduce challenging behaviors

Consistent use CAN improve behaviors/skills

Controversy

- Past iteration – positive and negative reinforcement (punishment for failure)
- Too intense for young children
- Neurodiversity (autism is a normal variation; "not-autistic")

Psychotherapies

Cognitive Behavioral Therapy

Thoughts, emotions, and behaviors
Helpful for co-occurring depression, anxiety

Adaptations

- Written, visual information
- Behavior >> Cognitive strategies
- Concrete language
- Psychoeducation about emotions

Dialectic Behavioral Therapy

Thoughts, emotions, and behaviors
+ mindfulness, emotion regulation, and distress tolerance

Helpful for suicidal ideation, self harm, destructive behaviors, emotional dysregulation

Pharmacologic Treatments

There are **NO** medications for the core symptoms of autism

Focus on treating co-occurring conditions

Medication selection follows similar algorithm as NT population

- Higher sensitivity to effects
- Increased likelihood of adverse effects
- Adjust titration schedule

Years of research, though inconsistent results in efficacy in treatment studies

- Varied etiology of "autism"

Pharmacologic Treatments

Serotonergic Agents

Regulate serotonin

- GI
- CV
- CNS

Theory: serotonin dysregulation → repetitive behaviors, anxiety, irritability, etc.

SSRI >>> SNRI, TCA

Atypical Antipsychotics

2 FDA approved for irritability associated with autism

- Risperidone
- Aripiprazole

Dopamine, serotonin, alpha-adrenergic, histaminergic receptors in CNS

Regular monitoring for side effects

Periodically re-evaluate need to continue treatment

Pharmacologic Treatments

Stimulants

Usually 1st line for co-occurring ADHD
PMH, FMH, PE (CV)

Side effects: decreased appetite, HTN, weight loss, sleep disruption, headache

Baseline sleep issues not necessarily predictive of stimulant related sleep issues

AMP – slightly more effective

MPH – usually better tolerated

Alpha-2-adrenergic Agonists

Clonidine and Guanfacine

Nonstimulant treatment for ADHD

Less effective than stimulants

Can be helpful with co-occurring sleep issues

Small study showing clonidine has positive effects

- Irritability
- Stereotypies
- Hyperactivity
- Hyperarousal

Social Supports

Board of Developmental Disabilities

School Based

Behavioral Support

Daily Living Skills

Social Skills Training/Groups

Sources

Ashworth, Runkumar, et al. "An Update on Psychopharmacological Treatment of Autism Spectrum Disorder." *Neurotherapeutics*, no. 1, Springer Science and Business Media LLC, Jan. 2022, pp. 248-62. Crossref, doi:10.1007/s13311-022-01185-1.

"Applied Behavior Analysis | Psychology Today." *Psychology Today*. <https://www.psychologytoday.com/us/therapy-types/applied-behavior-analysis>. Accessed 1 Mar. 2023.

Collins, Heidi, and Matthew Siegel. "Recognizing and Treating Comorbid Psychiatric Disorders in People With Autism." *Psychiatric Times*, Psychiatric Times, 29 Aug. 2019. <https://www.psychiatrictimes.com/news/recognizing-and-treating-comorbid-psychiatric-disorders-in-people-with-autism>.

Cooper, K., et al. "Adapting Psychological Therapies for Autism." *Research in Autism Spectrum Disorders*, Elsevier BV, Jan. 2018, pp. 43-50. Crossref, doi:10.1016/j.rasd.2017.11.002.

Edelson, Stephen M. "Understanding Challenging Behaviors in Autism Spectrum Disorder: A Multi-Component, Interdisciplinary Model." *Journal of Personalized Medicine*, no. 7, MDPI AG, July 2022, p. 1127. Crossref, doi:10.3390/jpm7071127.

Hartung, Anna, et al. "The Effect of Behavioral Therapy in Autism Spectrum Patients with Stolidity and/or Self-Destructive Behavior (DASD): Study Protocol for a Multicenter Randomized Controlled Trial." *BMC Psychiatry*, no. 1, Springer Science and Business Media LLC, Mar. 2020. Crossref, doi:10.1186/s12888-020-02031-4.

Leekam, Susan B., et al. "Restricted and Repetitive Behaviors in Autism Spectrum Disorders: A Review of Research in the Last Decade." *Psychological Bulletin*, no. 4, American Psychological Association (APA), 2011, pp. 342-69. Crossref, doi:10.1037/a0023341.

Pfiffner, Edith, et al. "Understanding Behavioral Rigidity in Autism Spectrum Conditions: The Role of Intentional Control." *Journal of Autism and Developmental Disorders*, Jan. 2017.

Roman, Maria, et al. "Psychiatric Comorbidities in Autism Spectrum Disorder: A Comparative Study Between DSM-IV-TR and DSM-5 Diagnosis." *International Journal of Clinical and Health Psychology*, May 2016.

Tian, Junbin, et al. "Repetitive Restricted Behaviors in Autism Spectrum Disorder: From Mechanism to Development of Therapeutics." *Frontiers in Neuroscience*, Frontiers Media SA, Mar. 2022. Crossref, doi:10.3389/fnins.2022.894677.

Williams, Diana, et al. "Problem Behaviors in Autism Spectrum Disorder: Association with Verbal Ability and Adapting/Doing Skills." *Journal of Autism and Developmental Disorders*, Nov. 2018.

Homes, Oliver D., et al. "Autism Spectrum Disorder: Consensus Guidelines on Assessment, Treatment and Research from the British Association for Psychopharmacology." *Journal of Psychopharmacology*, no. 1, SAGE Publications, Dec. 2017, pp. 3-20. Crossref, doi:10.1177/026981117741766.

MedNet21
Continued Learning and Professional Development

THE OHIO STATE UNIVERSITY
OHIO STATE MEDICAL CENTER